

UNIVERSIDADE DO ESTADO DE SANTA CATARINA – UDESC
CENTRO DE CIÊNCIAS AGROVETERINÁRIAS – CAV
PROGRAMA DE PÓS-GRADUAÇÃO EM CIÊNCIAS ANIMAL - PPGCA

ANDRÉ LUCIO FONTANA GOETTEN

**EFEITOS DOS ALCALÓIDES PIRROLIZIDÍNICOS PRESENTES EM *Senecio
brasiliensis* SOBRE OS GENES DA CASCATA PERIOVULATÓRIA EM CÉLULAS
DA GRANULOSA E SOBRE A PRODUÇÃO DE EMBRIÕES BOVINOS IN VITRO**

LAGES

2022

ANDRÉ LUCIO FONTANA GOETTEN

**EFEITOS DOS ALCALÓIDES PIRROLIZIDÍNICOS PRESENTES EM *Senecio
brasilensis* SOBRE OS GENES DA CASCATA PERIOVULATÓRIA EM CÉLULAS
DA GRANULOSA E SOBRE A PRODUÇÃO DE EMBRIÕES BOVINOS IN VITRO**

Tese apresentada ao Programa de Pós-graduação em Ciência Animal da Universidade do Estado de Santa Catarina, como requisito parcial para obtenção do título de doutor em Ciência Animal, área de concentração em Reprodução Animal.
Orientador: Prof. Dr. Alceu Mezzalana
Coorientador: Prof. Dr. Valério Valdetar Marques Portela Junior

LAGES

2022

Ficha catalográfica elaborada pelo programa de geração automática da Biblioteca Setorial do CAV/UDESC, com os dados fornecidos pelo(a) autor(a)

Goetten, André Lucio Fontana

Efeitos dos alcaloides pirrolizidínicos presentes em *Senecio brasiliensis* sobre os genes da cascata periovulatória em células da granulosa e sobre a produção de embriões bovinos in vitro / André Lucio Fontana Goetten. -- 2022.

106 p.

Orientador: Alceu Mezzalira

Coorientador: Valério Valdetar Marques Portela Junior

Tese (doutorado) -- Universidade do Estado de Santa Catarina, Centro de Ciências Agroveterinárias, Programa de Pós-Graduação em Ciência Animal, Lages, 2022.

1. Senecionina. 2. Seneciofilina. 3. Células da granulosa. 4. Ovários. 5. Embrião. I. Mezzalira, Alceu. II. Portela Junior, ValérioValdetar Marques. III. Universidade do Estado de Santa Catarina, Centro de Ciências Agroveterinárias, Programa de Pós-Graduação em Ciência Animal. IV. Título.

ANDRÉ LUCIO FONTANA GOETTEN

**EFEITOS DOS ALCALÓIDES PIRROLIZIDÍNICOS PRESENTES EM *Senecio
brasilensis* SOBRE OS GENES DA CASCATA PERIOVULATÓRIA EM CÉLULAS
DA GRANULOSA E SOBRE A PRODUÇÃO DE EMBRIÕES BOVINOS IN VITRO**

Tese apresentada ao Programa de Pós-graduação em Ciência Animal da Universidade do Estado de Santa Catarina, como requisito parcial para obtenção do título de doutor em Ciência Animal, área de concentração em Reprodução Animal.
Orientador: Prof. Dr. Alceu Mezzalana
Coorientador: Prof. Dr. Valério Valdetar Marques Portela Junior

BANCA EXAMINADORA

Coorientador:

Dr. Valério Valdetar Marques Portela Junior - UFSM

Membros:

Dr. Aldo Gava – UDESC

Dr. Maicon Gaissler Lorena Pinto - EPAGRI

Dr. Marcos Henrique Barreta – UFSC

Dr. Vitor Braga Rissi – UFSC

Lages, 28 de abril de 2022

Aos meus pais, Altamir (*in memoriam*) e Lela pelo exemplo de retidão e perseverança. A meus filhos, Áquila e Arthur, para mostrar-lhes que a busca pelo conhecimento deve ser incessante.

AGRADECIMENTOS

A Deus por estar sempre presente na minha vida e por ter colocado pessoas maravilhosas no meu caminho.

Ao Ir.: e amigo Professor Valério Portela, mentor e coorientador desse projeto, por ter sido o grande incentivador da retomada de minha vida acadêmica.

A minha esposa Lili pelo carinho, pelo apoio, pela compreensão nos momentos que não pude lhe dar a atenção merecida para dedicar-me a este projeto, mas principalmente por ter aceitado incondicionalmente adiar a realização de um grande sonho para que eu pudesse concluir meu doutorado.

Ao meu Orientador Prof. Alceu Mezzalira pela oportunidade concedida.

Aos colegas de LAFRA, Professor Marcos Henrique Barreta e Professor Vitor Braga Rissi pelo auxílio na condução dos experimentos e na resolução dos problemas encontrados.

Ao Biorep, especialmente às pós-graduandas Daniele Missio e Carolina Amaral, onde foram feitas parte das análises desse trabalho.

Aos Frigoríficos Ell'Golli e Verdi por terem gentilmente fornecido os ovários utilizados durante os experimentos.

A UFSC por ter, como parte de sua política para o desenvolvimento de pessoas, concedido o afastamento para minha capacitação.

Ao Curso de Medicina Veterinária e ao Programa de Pós-graduação em Ciência Animal do CAV/UDESC pelo ensino superior, mestrado e doutorado de excelência.

Aos Catarinenses, que com seus esforços e tributos viabilizam as atividades da UDESC.

“Explicar toda a natureza é uma tarefa difícil demais para qualquer homem ou qualquer época. É muito melhor fazer um pouco com certeza e deixar o resto para outros que vêm depois de você.” (ISAAC NEWTON)

RESUMO

As plantas contendo alcaloides pirrolizidínicos (APs) são amplamente distribuídas no mundo e provavelmente são as que mais causam intoxicação em animais de produção, animais silvestres e em humanos. No sul do Brasil o gênero *Senecio*, especialmente o *S. brasiliensis*, é responsável por mais de 50% das mortes de bovinos intoxicados por plantas. A intoxicação, que causa primariamente dano hepático, depende da biotransformação dos APs pela enzima citocromo P450 presente no fígado, mas que tem isoformas em outros órgãos. O curso da doença pode ser agudo ou crônico dependendo da frequência e da quantidade ingerida. Em vacas com seneciose crônica já foram descritas lesões ovarianas microscópicas semelhantes as lesões hepáticas e, em ovários suínos, uma isoforma de citocromo P450 foi identificada. A hipótese desse trabalho é que concentrações de APs insuficientes para causar seneciose atuam diretamente sobre as células ovarianas e/ou sobre os embriões, afetando a expressão dos genes da cascata ovulatória e o desenvolvimento embrionário inicial em bovinos. Os objetivos desse estudo foram avaliar se concentrações de alcaloides pirrolizidínicos, insuficientes para causar seneciose, atuam diretamente sobre: a) a expressão dos genes da cascata ovulatória nas células da granulosa bovinas, b) na expansão das células do cumulus-oócito e c) no desenvolvimento embrionário inicial em bovinos. Senecionina e seneciofilina, os dois PAs mais prevalentes no *S. brasiliensis*, foram testados nas concentrações de 0,03 μM e 0,3 μM em todos os experimentos. Células da granulosa foram cultivadas utilizando um modelo bem estabelecido no qual respondem ativamente ao LH in vitro. A concentração mais alta de ambos os APs reduziu a expressão de mRNA de *EREG*, *PTGS2* e *CTGF* 12 horas após o tratamento e de *AREG* as 24 horas após o tratamento. As concentrações de 0,03 μM senecionina as 12 horas, e 0,03 μM seneciofilina as 24 horas pós-tratamento também reduziram a expressão de *PTGS2*. Esses resultados sugerem que os APs podem prejudicar a ovulação. Utilizando um sistema de produção in vitro de embriões foram avaliadas: a expansão das células do complexo cumulus-oócito, a produção embrionária e a expressão de mRNA de genes marcadores de qualidade embrionária. A expansão do cumulus não foi impedida por nenhuma das doses de APs testadas. As taxas de clivagem e blastocisto também não diferiram do grupo controle quando os APs foram adicionados ao meio de maturação. Quando o tratamento ocorreu 18-20 horas após a inseminação, as taxas de clivagem e de blastocisto foram reduzidas por ambos os APs em todas as concentrações testadas. O risco de um suposto zigoto tratado com 0,03 μM de senecionina ou de seneciofilina chegar ao estágio de blastocisto foi reduzido em 29,7% e 38,5% comparados ao grupo não tratado, respectivamente. Para os grupos tratados com 0,3 μM de senecionina e seneciofilina esse risco foi reduzido em 54,1% e 47,3%, respectivamente. Em blastocistos, a seneciofilina reduziu a expressão o gene *TP53BP1* durante a maturação in vitro e aumentou a expressão do gene *RAD51* durante o cultivo in vitro, sugerindo que os APs poderiam determinar a quebra da fita dupla do DNA e a ativação de reparo via recombinação homóloga. A senecionina reduziu a expressão de mRNA de *GLUT3* e *BCL2*, sugerindo que os APs podem prejudicar o metabolismo da glicose e reduzir a atividade antiapoptótica em blastocistos bovinos. Em resumo, o presente estudo demonstra que a senecionina e a seneciofilina têm efeitos toxicológicos significativos em células da granulosa bovina, bem como em embriões bovinos produzidos in vitro, sugerindo que os PAs, mesmo em concentrações insuficientes para causar a seneciose, têm o potencial de afetar diretamente a função reprodutiva em bovinos.

Palavras-chave: senecionina, seneciofilina, células da granulosa, ovário, embrião.

ABSTRACT

Pyrrolizidine alkaloids (PAs)-containing plants are widely distributed in the world and are probably the most common poisonous plants affecting livestock, wildlife, and humans. In southern Brazil, the genus *Senecio*, especially *S. brasiliensis*, is responsible for more than 50% of the deaths of cattle poisoned by plants. Intoxication, which mainly causes liver damage, depends on the biotransformation of PAs by the enzyme cytochrome P450, which is present in the liver but has isoforms in other organs. The course of the disease can be acute or chronic, depending on the frequency and amount ingested. In cows with chronic seneciosis, microscopic ovarian lesions similar to liver lesions have already been described, in porcine ovaries, an isoform of cytochrome P450 has been identified. This work hypothesizes that concentrations of PAs insufficient to cause seneciosis act directly on ovarian cells and/or on embryos, affecting the expression of genes of the periovulatory cascade and early embryonic development in cattle. The objectives of this study were to evaluate whether concentrations of pyrrolizidine alkaloids, insufficient to cause seneciosis, act directly on: a) the expression of the genes of the periovulatory cascade in bovine granulosa cells, b) the expansion of cumulus-oocyte cells, and c) early embryonic development in cattle. Senecionine and seneciphylline, the two most prevalent PAs in *S. brasiliensis*, were tested at concentrations of 0.03 μM and 0.3 μM in all experiments. Granulosa cells were cultured using a well-established culture model in which they actively respond to LH. The higher concentration of both PAs reduced the mRNA expression of *EREG*, *PTGS2*, and *CTGF* 12 hours after treatment and *AREG* 24 hours after treatment. Concentrations of 0.03 μM senecionine at 12 hours and 0.03 μM seneciphylline at 24 hours post-treatment also reduced *PTGS2* expression. These results suggest that PAs can impair ovulation. Using the in vitro embryo production system, cumulus cell expansion, embryo production, and mRNA expression of embryonic quality marker genes were evaluated. Cumulus expansion was not impeded by any of the PAs tested. Cleavage and blastocyst rates also did not differ from the control group when PAs were added to the maturation medium. When treatment was added 18-20 hours post-insemination, cleavage and blastocyst rates were reduced by both PAs at all concentrations tested. The risk of a putative zygote treated with 0.03 μM senecionine or seneciphylline reaching the blastocyst stage was reduced by 29.7% and 38.5% compared to the untreated group, respectively. For the groups treated with 0.3 μM of senecionine and seneciphylline, this risk was reduced by 54.1% and 47.3%, respectively. In blastocysts, seneciphylline downregulated the *TP53BP1* gene during in vitro maturation and the *RAD51* gene was upregulated during in vitro culture, suggesting that PAs could determine DNA double-strand breaks and activation of the homolog recombination repair pathway. Senecionine reduced mRNA expression of *GLUT3* and *BCL2*, suggesting that PAs may impair glucose metabolism and reduce antiapoptotic activity in bovine blastocysts. In summary, the present study demonstrates that senecionine and seneciphylline have significant toxicological effects on bovine granulosa cells as well as in vitro produced bovine embryos, suggesting that PAs, even at concentrations insufficient to cause seneciosis, have the potential to directly affect the reproductive function in cattle.

Keywords: senecionine, seneciphylline, granulosa cells, ovary, embryo.

LISTA DE ILUSTRAÇÕES

Figura 1 – Representação gráfica da comunicação bidirecional entre o oócito e as células do cumulus	25
Figura 2 – Quatro abordagens para o reparo das quebras na fita dupla do DNA.....	31

ARTIGO I

Figure 1 – Effect of Senecionine (A) or Seneciophylline (B) on expression of periovulatory EGF-Like cascade genes in bovine granulosa cells.....	47
Figure 2 – Effect of graded concentrations (0, 0.03, and 0.3 μ M) of A) Senecionine or B) Seneciophylline on cumulus cells expansion.	48
Figure 3 – Risk ratio for cleavage on day 2 (A; B) or blastocyst on day 7 (C; D) after in vitro fertilization of COCs treated with graded concentrations of pyrrolizidine alkaloids during in vitro maturation.	49
Figure 4 – Relative mRNA expression of genes associated with DNA double-strand repair in bovine embryos produced in vitro and treated with graded concentrations of pyrrolizidine alkaloids during in vitro maturation.....	50
Figure 5 – Relative mRNA expression of BAX, BCL2 and BAX/BCL2 ratio in bovine embryos produced in vitro and treated with graded concentrations of pyrrolizidine alkaloids during in vitro maturation.	51
Figure 6 – Relative mRNA expression of GLUT1 and GLUT3 in bovine embryos produced in vitro and treated with graded concentrations of pyrrolizidine alkaloids during in vitro maturation.	52
Figure 7 – Risk ratio for cleavage on day 2 (A; B) or blastocyst on day 7 (C; D) after in vitro fertilization of zygotes treated with graded concentrations of pyrrolizidine alkaloids added to the medium at 18-20 hpi.....	53
Figure 8 – Relative mRNA expression of genes associated with DNA double-strand repair in bovine in vitro produced blastocysts treated with graded concentrations of pyrrolizidine alkaloids added to the medium at 18-20 hpi.	54
Figure 9 – Relative mRNA expression of BAX, BCL2, and bax/bcl2 ratio in bovine in vitro produced blastocysts treated with graded concentrations of pyrrolizidine alkaloids added to the medium at 18-20 hpi.	55

Figure 10– Relative mRNA expression of GLUT1 and GLUT3 in bovine in vitro produced blastocysts treated with graded concentrations of pyrrolizidine alkaloids added to the medium at 18-20 hpi.	56
---	----

ARTIGO II

Fig. 1 - Abundance of A) Prostaglandin-endoperoxide Synthase 2 (PTGS2), B) Epiregulin (EREG), C) Hyaluronan Synthase 2 (HAS2) and D) TNF Alpha Induced Protein 6 (TNFAIP6) transcripts in bovine cumulus cells from COCs submitted to IVM for 3, 6, 12 or 24 hours in medium supplemented with bovine serum albumin (BSA) or fetal calf serum (FCS).	91
Fig. 2 – Abundance of A) X-ray repair cross complementing 6 (XRCC6), B) X-ray repair cross complementing 5 (XRCC5), C) Protein kinase DNA-activated, catalytic subunit (DNAPK), D) growth arrest and DNA damage inducible beta (GADD45B) and E) tumor protein p53 binding protein 1 (TP53BP1) transcripts in bovine cumulus cells from COCs submitted to IVM for 3, 6, 12 or 24 hours in medium supplemented with bovine serum albumin (BSA) or fetal calf serum (FCS)	92
Fig. 3 – Abundance of A) RAD50 double strand break repair protein (RAD50), B) RAD51 recombinase (RAD51), C) RAD52 homolog DNA repair protein (RAD52), D) ATM serine/threonine kinase (ATM), E) BRCA1 DNA repair associated (BRCA1) and F) BRCA2 DNA repair associated (BRCA2) transcripts in bovine cumulus cells from COCs submitted to IVM for 3, 6, 12 or 24 hours in medium supplemented with bovine serum albumin (BSA) or fetal calf serum (FCS).	93

LISTA DE TABELAS

ARTIGO I

Table 1 – Details of primers used for gene expression analysis RT-qPCR.....	42
Table 2 – Cleavage rates at 48 hpi and blastocyst rates at 168 hpi of COCs treated with graded concentrations of pyrrolizidine alkaloids during in vitro maturation.	48
Table 3 – Cleavage rates at 48 hpi and blastocyst rates at 168 hpi of COCs treated with graded concentrations of pyrrolizidine alkaloids added to IVC medium.	52

ARTIGO II

Table 1 - Information of specific primers used for amplification in real-time PCR.	90
---	----

LISTA DE ABREVIATURAS E SIGLAS

AP	Alcaloides pirrolizidínicos
AREG	Anfiregulina
ATM	<i>Ataxia-telangiectasia mutated</i>
ATR	<i>ATM and RAD3 related</i>
BMP15	Proteína Morfogenética Óssea 15
BRCA1	<i>Breast Cancer 1 DNA repair associated</i>
BRCA2	<i>Breast Cancer 2 DNA repair associated</i>
BSA	Albumina Sérica Bovina
cAMP / AMPc	Monofosfato de Adenosina Cíclico
CCO / COC	Complexo cumulus-oócito
cGMP / GMPc	Monofosfato de Guanosina Cíclico
CI	<i>Intervalo de confiança</i>
CTIP	Proteína <i>CtBP-interacting</i>
DDR	Resposta ao dano no DNA
DNAPK	Proteína Quinase dependente de DNA
DNAPK-cs	Proteína Quinase dependente de DNA, subunidade catalítica
DSB	Quebra de fita dupla no DNA
EGF	Fator de Crescimento Epidermal
EGF-Like	Fator de Crescimento semelhante ao EGF
EGFR	Receptor de EGF
EREG	Epiregulina
FCS	Soro Fetal Bovino
FGF	Fatores de Crescimento Fibroblásticos
FSH	Hormônio Folículo Estimulante
GADD45B	<i>Growth Arrest and DNA Damage Inducible Protein, Beta</i>
GDF9	Fator de Diferenciação de Crescimento 9

H2AFZ	Família de histonas H2A, membro Z
HAS2	Hialurona Sintetase 2
HPI	Horas pós inseminação
HR	Recombinação Homóloga
IVM	Maturação <i>in vitro</i>
IVP	Produção de embriões <i>in vitro</i>
LH	Hormônio Luteinizante
LHR	Receptor de Hormônio Luteinizante
LIG4	DNA ligase IV
MMEJ	União Terminal Mediada por Micro-homologia
MRN	Complexo Mre11-Rad50-Nbs1
NHEJ	União Terminal Não Homóloga
PARP	Poli(ADP-ribose) polimerase
PDE	Fosfodiesterase
PGE2	Prostaglandina E2
PIKK	Proteína quinase semelhante a fosfatidilinositol-3 quinase
PPIB	Ciclofilina B
PTGS2	Prostaglandina Sintetase 2
PTIP	<i>PAX-interacting protein 1</i>
PTX3	Pentraxina 3
PV	Peso Vivo
RAD50	<i>DNA repair protein RAD50</i>
RAD51	<i>DNA repair protein RAD51</i>
RAD52	<i>DNA repair protein RAD52</i>
RIF1	<i>Rap1-interacting factor 1</i>
RPA	Proteína de Replicação A
S.E.M.	Erro padrão da média
SSA	Anelamento de Fitas Simples

SSB	Quebra de fita simples no DNA
ssDNA	Fita Simples de DNA
TBP	Proteína de Ligação TATA
TNAFAIP6	Proteína Indutora do Fator de Necrose Tumoral 6
TP53BP1	Proteína de Ligação de Proteína Tumoral P53 1
XRCC4	Proteína 4 de complementação cruzada de reparo de raios X
XRCC5	Proteína 5 de complementação cruzada de reparo de raios-X
XRCC6	Proteína 6 de complementação cruzada de reparo de raios-X
YAP	<i>Yes-Associated Protein</i>

LISTA DE SÍMBOLOS

%	Porcento
°C	Graus Celsius
H	Horas
IU	Unidades Internacionais
mL	Mililitro
µg	Micrograma
Mm	Milímetro
µL	Microlitro
CO ₂	Dióxido de Carbono
Ng	Nanograma
µM	Micromolar

SUMÁRIO

1 INTRODUÇÃO.....	15
2 REVISÃO DE LITERATURA	19
2.1 <i>SENECIO BRASILIENSIS</i> E SEUS EFEITOS NOS BOVINOS	19
2.2 A IMPORTÂNCIA DA COMUNICAÇÃO ENTRE AS CÉLULAS DO COMPLEXO CUMULUS-OÓCITO	23
2.3 A EXPANSÃO DAS CÉLULAS DO CUMULUS	26
2.4 VIAS DE SINALIZAÇÃO E REPARO DOS DANOS DE DNA	27
2.4.1 Reparos de quebras na fita dupla do DNA	30
2.4.1.1 <i>União terminal não homóloga (NHEJ)</i>	32
2.4.1.2 <i>Recombinação homóloga (HR)</i>	33
3 ARTIGO I - <i>Senecio brasiliensis</i>-DERIVED PYRROLIZIDINE ALKALOIDS DECREASE OVULATORY CASCADE GENE EXPRESSION AND BOVINE EMBRYOS PRODUCTION.....	36
3.1 INTRODUCTION.....	38
3.2 MATERIAL AND METHODS	40
3.2.1 Granulosa cell culture.....	40
3.2.2 Cumulus-oocyte complexes recovery and in vitro maturation (IVM)	41
3.2.3 In vitro fertilization (IVF) and in vitro embryo culture (IVC).....	41
3.2.4 RNA extraction, reverse transcription, and real-time polymerase chain reaction .	42
3.3 STATISTICAL ANALYSIS	44
3.3.1 Experimental design	44
3.3.1.1 <i>Experiment 1 – Effects of PAs on the periovulatory EGF-Like cascade genes in granulosa cells.....</i>	44
3.3.1.2 <i>Experiment 2 - Effect of PAs on cumulus-oocyte complexes expansion.....</i>	45
3.3.1.3 <i>Experiment 3 - Effects of PAs added to the IVM medium on bovine embryo production and quality.</i>	45
3.3.1.4 <i>Experiment 4 - Effects of PAs added to the IVC medium on bovine embryo production and quality.</i>	46

3.4 RESULTS.....	46
3.4.1 mRNA relative expression of periovulatory EGF-Like cascade genes in bovine granulosa cells exposed to PAs.....	46
3.4.2 PAs have no effect on cumulus cell expansion in in vitro matured COCs.....	48
3.4.3 Effects of PAs added to the IVM medium on bovine embryo production and quality.....	48
3.4.4 Effects of PAs added to the IVC medium on bovine embryo production and quality.....	52
3.5 DISCUSSION	56
3.6 REFERENCES.....	60
4 ARTIGO II – EXPRESSION PROFILE OF KEY GENES INVOLVED IN DNA REPAIR MECHANISMS IN BOVINE CUMULUS CELLS CULTURED WITH BOVINE SERUM ALBUMIN OR FETAL CALF SERUM	66
4.1 ABSTRACT	68
4.2 INTRODUCTION.....	68
4.3 MATERIAL AND METHODS	72
4.3.1 In vitro maturation of Cumulus-oocyte complexes.....	72
4.3.2 Nucleic acid extraction and qPCR.....	73
4.3.3 Statistical analysis	74
4.4 RESULTS.....	75
4.5 DISCUSSION	76
4.6 REFERENCES.....	81
REFERÊNCIAS	94

1 INTRODUÇÃO

A intoxicação por plantas ainda hoje é uma das causas mais comuns de morte de animais de produção adultos, determinando expressivas perdas econômicas, diretas e indiretas, no sistema produtivo pecuário. Contudo, a avaliação precisa do prejuízo decorrente das intoxicações por plantas não é tarefa fácil devido a escassez de informações confiáveis já que nem todos os surtos são notificados e, principalmente, por conta da falta de dados a respeito dos casos nos quais a intoxicação não provoca a morte imediata, mas determina queda no desempenho dos animais.

Falhas de manejo nas áreas de pastagem ou de lavouras favorecem o aparecimento de plantas invasoras, muitas das quais tem potencial tóxico para várias espécies animais, inclusive o homem. Das plantas invasoras do sul do Brasil, aquelas pertencentes ao gênero *Senecio* spp. são as que determinam as maiores perdas econômicas na criação de bovinos, dentre elas o *Senecio brasiliensis* tem lugar de destaque. Por ser altamente resistente e com grande capacidade de adaptação às condições edafoclimáticas de onde se estabelece, o *S. brasiliensis* é uma planta amplamente difundida pela América do Sul e em especial, na região sul do Brasil (PEDROSO et al., 2007; RISSI et al., 2007; GRECCO et al., 2010). Nesta região a maioria dos bovinos é criada em sistemas extensivos o que facilita o aparecimento dos quadros de intoxicação por *Senecio* spp., especialmente em épocas em que a pastagem é escassa. Entretanto a seneciose, intoxicação por plantas do gênero *Senecio*, também tem sido descrita em sistemas intensivos e semi-intensivos quando os animais ingerem a planta involuntariamente seja através de volumoso contaminado ou pelo hábito de pastejo particular dos bovinos que dificulta a seleção do alimento durante o bocado (KARAM et al., 2004; BIFFI et al., 2019).

Os princípios ativos tóxicos associados ao *S. brasiliensis* pertencem ao grupo dos alcaloides pirrolizidínicos (APs) que, quando biotransformados em pirróis pelas enzimas do citocromo P-450, são capazes alterar a fisiologia celular, especialmente dos hepatócitos, já que causam lesões irreversíveis nessas células. Já se sabe que uma parte dos pirróis produzidos no fígado pode atingir outros órgãos através da circulação sistêmica e causar lesão extra-hepática. As lesões extra-hepáticas também podem ocorrer pela produção de pirróis localmente, já que os complexos do citocromo P-450 são encontrados em outros órgãos (PRAKASH et al., 1999).

A redução das taxas reprodutivas de fêmeas bovinas acometidas por seneciose sempre foi relacionada ao declínio do status nutricional em decorrência da lesão hepática nesses animais. Contudo, há indícios de atuação dos APs diretamente em células ovarianas já que ovários de suínos são capazes de expressar e transcrever a enzima citocromo P450 e suas isoformas e ovários de bovinos com seneciose crônica apresentaram, em células luteais, lesões semelhantes aquelas presentes nos hepatócitos. Também há relatos de que a exposição pré-natal a APs pode prejudicar o desenvolvimento físico e comportamental em ratos (LEIGHTON et al., 1995; SANDINI et al., 2014; KRABBE et al., 2015). Quando associamos problemas de infertilidade sem causas aparentes em bovinos a essas evidências de atuação dos APs sobre órgãos reprodutivos, surge a hipótese de que, mesmo em concentrações insuficientes para causar a seneciose crônica, os APs atuam diretamente sobre as células ovarianas e/ou sobre os embriões, afetando a expressão dos genes da cascata ovulatória e o desenvolvimento embrionário inicial em bovinos. Os objetivos do primeiro estudo foram avaliar se concentrações de alcaloides pirrolizidínicos, insuficientes para causar seneciose, atuam diretamente sobre: a) a expressão dos genes da cascata ovulatória nas células da granulosa bovinas, b) na expansão das células do complexo cumulus-oócito e c) no desenvolvimento embrionário inicial em bovinos.

Essa tese contém um segundo artigo que trata do padrão da expressão de genes envolvidos no reparo da fita dupla do DNA em células do cumulus bovinas cultivadas sob influência de soro fetal bovino (FCS) ou de albumina sérica bovina (BSA), cujos resultados foram utilizados para a definição da inclusão de FCS durante a maturação in vitro nos experimentos que testaram os efeitos dos APs na produção in vitro de embriões.

Apesar de seu uso generalizado, a eficácia da maturação in vitro (IVM) é menor que a observada em oócitos maturados in vivo. Portanto, informações mais detalhadas sobre as vias moleculares que controlam a expansão das células do cumulus e a maturação do oócito são necessárias para melhorar os protocolos de IVM (CAIXETA et al., 2013a; DE VOS et al., 2021). A regulação da maturação do oócito e da expansão das células do cumulus depende da comunicação entre esses dois compartimentos do complexo cumulus-oócito (CCO) e envolve trocas bidirecionais por meio de junções comunicantes e sinalização parácrina. As junções comunicantes dos CCOs permitem a troca de pequenas moléculas que dão suporte nutricional e regulam a progressão meiótica do oócito o qual, por sua vez, secreta fatores que modulam a proliferação, expressão gênica e função das células do cumulus (TANGHE et al., 2002; SUGIMURA; RICHANI; GILCHRIST, 2018).

As condições de cultura durante a produção de embriões in vitro (PIV) podem afetar os padrões de expressão gênica e, conseqüentemente, o potencial de desenvolvimento embrionário, indicando que o ambiente adverso gerado pelas condições in vitro prejudica a comunicação bidirecional dos CCOs. Assim, diversas alterações nas condições ou nos meios de cultura foram propostas na expectativa de minimizar essas adversidades. A escolha da fonte de suplementação proteica no meio de maturação, por exemplo, tem efeitos profundos nas taxas de desenvolvimento embrionário, justificando o uso de soro durante a IVM na maioria dos sistemas. No entanto, o uso de soro apresenta desvantagens científicas, sanitárias e éticas determinando a busca de outras fontes proteicas que possam substituí-lo e a albumina sérica bovina é a fonte proteica mais utilizada para reposição sérica. Infelizmente, a ausência de soro no meio IVM resulta em redução da taxa de blastocisto, menor grau de expansão do cumulus e maior fragmentação de DNA nas células do cumulus (LONERGAN et al., 2003; VAN DER VALK et al., 2018).

Entre os diferentes tipos de danos ao DNA induzidos por genotoxinas ambientais, as quebras de fita dupla do DNA (DSB) são o tipo de dano mais deletério. Como CCOs maturados in vitro em meio sem soro apresentam alta taxa de apoptose e fragmentação de DNA nas células do cumulus os mecanismos de reparo das DSB podem ser comprometidos nessas condições de cultura celular. Esse reparo é coordenado principalmente por duas vias moleculares: recombinação homóloga (HR) e união terminal não homóloga (NHEJ). Apesar da presença de genes codificadores de mRNA implicados nesses processos ter sido demonstrada em oócitos de várias espécies, incluindo os bovinos, e em células do cumulus em humanos e ratos, não há informações sobre a expressão de genes envolvidos nas vias HR e NHEJ nas células do cumulus em bovinos (PANIER; BOULTON, 2014; SUN et al., 2018).

Considerando que a interação bilateral coordenada entre as células do cumulus e o oócito é essencial para a aquisição da plena capacidade de ambos os tipos celulares em promover a embriogênese inicial, as informações armazenadas nas células do cumulus podem ser marcadores diretos para o futuro desenvolvimento do oócito. Portanto, os padrões de expressão gênica em células do cumulus biopsiadas podem servir como marcadores não invasivos para avaliar a maturação do oócito. É por isso que o perfil de expressão de genes de reparo de DNA em células do cumulus bovino é uma questão intrigante que pode fornecer conhecimento para melhorar a eficiência da IVM. Assim, os objetivos desse segundo estudo foram: (1) caracterizar o perfil de expressão de transcritos de genes selecionados envolvidos nos mecanismos de reparo das quebras de fita dupla de DNA em células do cumulus bovinas

obtidas a partir de COCs maturados in vitro; e (2) investigar se a suplementação do meio de maturação in vitro com BSA ou FCS afeta a expressão da transcrição desses genes.

2 REVISÃO DE LITERATURA

2.1 *Senecio brasiliensis* E SEUS EFEITOS NOS BOVINOS

Já foram descritas mais de 1200 espécies do gênero *Senecio* espalhadas por praticamente todo o planeta, exceto nas regiões polares e na região Amazônica. De todas as espécies identificadas apenas cerca de 25 a 30 demonstraram ser tóxicas para animais domésticos (TOKARNIA; DOBEREINER; PEIXOTO, 2012). No Brasil, o número de espécies do gênero *Senecio* é divergente, variando entre 67 e 128 de acordo com o critério adotado para a classificação (HIND, 1993; MÉNDEZ; RIET-CORREA, 2008). *Senecio brasiliensis* é a espécie mais abundante no Brasil, amplamente distribuída no Sul e em pequenas áreas altas e frias da região Sudeste do país (PEDROSO et al., 2007; RISSI et al., 2007; GRECCO et al., 2010). Esta planta apresenta características adaptativas tais como a dispersão de sementes pelo vento, a independência de polinizador específico, a competição por alelopatia e a capacidade de crescimento mesmo em solos inférteis as quais lhe conferem grande facilidade de propagação, principalmente sob condições climáticas ideais (DANA et al., 2021).

S. brasiliensis é uma planta invasora de pastagens, mas de pouca palatabilidade. Portanto, é consumida naturalmente pelos bovinos somente sob determinadas condições, normalmente associadas à escassez de alimentos (KARAM et al., 2004; BARROS et al., 2007; MÉNDEZ; RIET-CORREA, 2008; TOKARNIA; DOBEREINER; PEIXOTO, 2012). A ingestão ocorre principalmente a partir da segunda metade do outono até o final do inverno, quando diminui muito a disponibilidade de pastagem (BASILE et al., 2005; BARROS et al., 2007). Contudo, sob influência de precipitação, umidade do solo e cobertura vegetal favoráveis à emergência de espécies de *Senecio*, o seu desenvolvimento pode ocorrer em qualquer época do ano e, conseqüentemente, a ingestão e a intoxicação podem se manifestar durante todo o ano (KARAM et al., 2004). A superlotação de bovinos e a alta infestação de *Senecio* spp. nas pastagens favorecem a ingestão da planta e, se plantas novas estão estreitamente associadas ao capim, o perigo de ingestão pelos bovinos é ainda maior em função da pouca seletividade de pastejo nessa espécie (KARAM et al., 2004; RISSI et al., 2007; GRECCO et al., 2010; GIARETTA et al., 2014). Os animais também podem ser

intoxicados pela ingestão involuntária da planta quando são alimentados com feno e silagem contaminados (BASILE et al., 2005; BIFFI et al., 2019).

Alguns surtos de intoxicação por *Senecio* spp. foram descritos em bovinos jovens (PRAKASH et al., 1999; BASILE et al., 2005) mas, em geral, a Seneciose é observada em animais adultos (CHEEKE, 1988) por ingerirem, ao longo do tempo, maior quantidade de *Senecio* spp. já que permanecem por um maior tempo na propriedade (GORDON; COLEMAN; GRISHAM, 2000; GIARETTA et al., 2014). A morbidade é variável, entre 1 e 30%, e a letalidade é de praticamente 100% (GORDON; COLEMAN; GRISHAM, 2000).

O princípio tóxico presente nas espécies de *Senecio* pertence ao grupo dos alcaloides pirrolizidínicos cujos representantes identificados em plantas de *S. brasiliensis* são senecionina, seneciofilina, integerrimina, retrorsina, usaramina e jacobina (HARTMANN; WITTE, 1995; TRIGO et al., 2003). Destes, a senecionina e a seneciofilina são os mais prevalentes (ADAMS; GIANTURCO, 1956). Quimicamente os APs são ésteres de aminoálcoois com um núcleo pirrolizidínico (necina) e ácidos alifáticos (ácidos nélicos). A característica que determina a capacidade de um AP ser tóxico é a presença de dupla ligação entre os carbonos C1 e C2 da necina, já que aqueles que possuem a necina saturada não são tóxicos aos mamíferos (SILVA; BOLZAN; HEINZMANN, 2006).

O teor de APs varia entre as diferentes espécies de plantas e, embora sejam sintetizados nas raízes, existe variação entre as partes de uma mesma planta. O conteúdo de APs em espécies do gênero *Senecio* spp. varia muito durante o ciclo de crescimento da planta e de ano para ano (SANDINI; SAYURI; BERTO, 2013). Em plantas de *S. brasiliensis* o teor de APs apresenta variação sazonal, atingindo a máxima concentração nos meses de inverno (KARAM et al., 2004).

APs são compostos estáveis que, por si só, não apresentam toxicidade até que sejam biotransformados (BARROS et al., 2007). Após serem absorvidos no intestino, os APs são transportados até o fígado onde são primariamente metabolizados pelas enzimas do sistema citocromo P-450, originando pirróis (BARROS et al., 2007; TOKARNIA; DOBEREINER; PEIXOTO, 2012). Os pirróis são compostos altamente tóxicos, reativos e alquilantes que se ligam facilmente às moléculas proteicas e de ácidos nucleicos. No fígado atuam inibindo a mitose, promovem necrose e redução do número de hepatócitos, dando início ao quadro clínico e a morte devido à disfunção hepática (PRAKASH et al., 1999). Foi proposto que a inibição da mitose poderia ser a causa da megalocitose que precede a necrose de hepatócitos nos casos de seneciose. A inibição da citocinese sem a interrupção da síntese de DNA no

núcleo dessas células são a base desse mecanismo (SKILLETER; MATTOCKS; NEAL, 1988). Contudo, Prakash et al. (1999) sugeriram que essa megolocitose é causada por alterações no ciclo celular devido a danos no DNA dos hepatócitos.

Os efeitos hepatotóxicos dos pirróis são progressivos e irreversíveis (CHEEKE, 1988), porém quando os animais ingerem pequenas quantidades diárias de *S. brasiliensis* durante um tempo prolongado, um período de latência de várias semanas e até meses pode ocorrer até que sejam observados os primeiros sinais clínicos (KARAM et al., 2002; BASILE et al., 2005), simulando um efeito cumulativo desses xenobióticos (TOKARNIA; DOBEREINER, 1984). Quando o diagnóstico de intoxicação por *Senecio* spp. é confirmado em um rebanho bovino, provavelmente vários animais já estejam subcl clinicamente afetados, isto é, apresentam lesões hepáticas, mas ainda sem apresentar sinais clínicos. Como as lesões causadas pela planta são progressivas, as mortes desses bovinos subcl clinicamente afetados podem ocorrer bastante tempo após a ingestão da planta (MOLYNEUX; JOHNSON; STUART, 1988; TRIGO et al., 2003; CHENG et al., 2011). Existem relatos de animais que tiveram diagnóstico de lesão hepática decorrente da ingestão de *S. brasiliensis* e que depois de transcorridos mais de dois anos, tempo suficiente para duas gestações de uma fêmea bovina, não haviam sequer apresentado sinais clínicos (GUTIÉRREZ; CAMPBELL; WEBB, 1997), porém, o desempenho produtivo e reprodutivo desses animais poderia estar comprometido nesse período pois existe possibilidade de toxicidade fetal após ingestão materna de *Senecio* spp. (MOLYNEUX et al., 2011).

Tanto a forma aguda quanto a forma crônica de intoxicação por *S. brasiliensis* em bovinos foram demonstradas experimentalmente. Uma única administração das partes aéreas da planta verde fresca, correspondente a 17,5 e 35 g/kg de peso vivo (PV), foi capaz de reproduzir a seneciose aguda. Doses mais baixas, de 5-10 g/kg PV, também administradas somente uma vez ou doses diárias de 0,625-5 g/kg PV, totalizando 75-150g/kg PV, produziram a forma crônica da doença. A manifestação crônica da seneciose também foi observada após administração semanal das doses de 2,18-8,75 g/kg PV, totalizando 61,25-78,75 g/kg PV, durante o período de 1 a 8 meses (TOKARNIA; DOBEREINER, 1984). Teores de seneciofilina maiores que 0,3 µM foram encontrados no sangue e no leite de bovinos durante as primeiras 18 horas depois que receberam 1 mg/kg do alcaloide por via oral. Depois de 54 horas, níveis sanguíneos de 0,033 µM ainda estavam presentes (CANDRIAN et al., 1991). Considerando que os teores dos APs em *S. brasiliensis* variaram entre 0,178% a 0,31% (KARAM et al., 2004; LUCENA et al., 2010), a ingestão de 0,3 – 0,5

g/kg PV das partes aéreas da planta, quantidade menor que a dose mínima que causar a seneciose crônica, seria suficiente para determinar níveis sanguíneos de seneciofilina equivalentes a 0.3 μ M.

Embora a sensibilidade aos APs seja bastante variável entre espécies e entre os diferentes tipos de tecidos nos mamíferos, os APs in vitro determinaram efeitos citotóxicos e prejudicaram via de síntese de prostanóides em células do endotélio umbilical humano (HUVEC) em doses bem maiores que as que produziram sintomas em bovinos experimentalmente intoxicados (EBMEYER et al., 2019). Da mesma forma, a dose experimental suficiente para destruir 50% dos hepatócitos primários e células endoteliais sinusoidais do fígado (LSECs) de camundongo (HESSEL-PRAS et al., 2020) foi maior que a menor dose tóxica capaz de induzir experimentalmente a seneciose em bovinos. Além disso, a citotoxicidade em ambos os modelos experimentais só se manifestou após a bioativação dos APs, sem a qual somente níveis tão altos quanto 500 μ M de senecionina prejudicaram as LSECs. Por sua vez nenhum efeito foi observado em HUVECs para os APs equimidina, heliotrina, lasiocarpina, senecionina, senkirkina e platifilina na ausência de um sistema de metabolização externo, até a concentração mais alta testada de 500 μ M.

Os prejuízos econômicos decorrentes de intoxicações por plantas nos rebanhos brasileiros são de difícil determinação em função dos poucos dados disponíveis a esse respeito, contudo estima-se que no mínimo 5% da população bovina morrem anualmente no Brasil (PESSOA; MEDEIROS; RIET-CORREA, 2013) e dados de laboratórios de diagnóstico mostram que entre 7,5 e 15,83% desses casos devem-se à intoxicação por plantas (RIET-CORREA; MEDEIROS, 2001; PEDROSO et al., 2007; RISSI et al., 2007). A ingestão de plantas do gênero *Senecio*, em especial de *S. brasiliensis*, é considerada a principal intoxicação por plantas e uma das principais causas de morte entre bovinos na região Sul do Brasil, respondendo por cerca de 50 – 60% dos óbitos decorrentes de intoxicações nessa espécie (KARAM; SCHILD; MELLO, 2011).

Segundo o IBGE (2020) a população de bovinos na região Sul do Brasil é de pouco mais de 24 milhões de cabeças. Considerando o valor médio de US\$ 250,00 por animal pode-se inferir que as perdas anuais decorrentes de mortes causadas pela Seneciose em bovinos, apenas nos três Estados do Sul do país, possam passar de US\$ 25 milhões. Presume-se que estas perdas sejam ainda maiores já que animais com intoxicação crônica e que ainda não manifestem sinais clínicos da Seneciose possam apresentar baixa taxa de produção de carne e

leite, bem como subfertilidade, em função dos danos hepáticos que prejudicam o metabolismo animal (GÓRNIAK, 2008).

Apesar de o tecido hepático ser o principal alvo das lesões causadas pela ingestão de *S. brasiliensis*, outros tecidos e órgãos, como pulmões e rins podem sofrer ação dos AP uma vez que as enzimas do citocromo P-450 também estão presentes nesses locais (PRAKASH et al., 1999). Corroborando essa informação, já foi demonstrado que alguns APs podem ser metabolizados pela isoforma CYP1A1 do citocromo P-450 presente em astrócitos de camundongos (NASCIMENTO et al., 2017). Com base nos efeitos dos APs em múltiplos tecidos foi sugerido que essa ação deletéria também seja capaz de causar danos diretos ao sistema reprodutivo feminino, bem como afetar o desenvolvimento embrionário inicial, uma vez que já foram descritos megalocitose em células luteais bovinas causada pela Seneciose (KRABBE et al., 2015), expressão do Citocromo P-450 1A1 em células da granulosa de suínos (LEIGHTON et al., 1995) e atraso no desenvolvimento físico e comportamental em ratos recém nascidos que tiveram exposição pré-natal a APs (SANDINI et al., 2014).

Boa parte dos casos de infertilidade em bovinos não tem etiologia estabelecida ou são erroneamente atribuídos a agentes infecciosos (ANTONIASSI et al., 2013), porém o desconhecimento dos mecanismos patogénéticos das fitotoxinas sobre a reprodução faz com que sua importância não seja completamente reconhecida (KRABBE et al., 2015). Além disso, até o momento, não existem informações a respeito dos efeitos de APs sobre a fisiologia ovariana ou sobre o desenvolvimento inicial em bovinos. Dessa forma, o estudo das possíveis alterações reprodutivas causadas diretamente por APs pode contribuir para elucidar a causa de uma fração dos casos de infertilidade sem causa aparente em bovinos.

2.2 A IMPORTÂNCIA DA COMUNICAÇÃO ENTRE AS CÉLULAS DO COMPLEXO CUMULUS-OÓCITO

O complexo cumulus-oócito (CCO) pode ser definido como um conjunto de células da granulosa intimamente associadas, denominadas células do cumulus, circundando o oócito em um folículo antral. Além desses dois tipos celulares, as células da granulosa mural e da teca que também compõem o folículo, desempenham suas funções específicas e interligadas para garantir que o oócito tenha atingido seu total desenvolvimento e esteja apto para ser fecundado no momento da ovulação (TANGHE et al., 2002; SUGIMURA; RICHANI; GILCHRIST, 2018). Nesse sistema, os oócitos regulam a proliferação, expressão gênica e

função das células do cumulus, enquanto as células do cumulus fornecem o suporte nutricional essencial para o crescimento e desenvolvimento do oócito (SUGIURA et al., 2008).

No CCO, o oócito e as células do cumulus estão conectados por junções comunicantes (“*gap junctions*”). Esse tipo de conexão também acontece com as células do cumulus entre si e com as células da granulosa mural. Canais de membrana intercelulares, chamados projeções transzonais, estão associados a essas junções comunicantes e permitem a transferência física de pequenos fatores reguladores e metabólitos entre o oócito, as células do cumulus e as células murais da granulosa. Além disso, o oócito secreta fatores parácrinos capazes de regular várias funções biológicas das células somáticas que o rodeiam, configurando um sistema de comunicação bidirecional dinâmico entre duas categorias de células distintas (RUSSELL et al., 2016).

A importância dessas trocas bidirecionais ficou comprovada quando oócitos bovinos desnudos maturados *in vitro* ou na presença de inibidores de junções comunicantes apresentaram alterações metabólicas e menor capacidade de sustentar a fertilização e o subsequente desenvolvimento embrionário, comparados com CCOs maturados nas mesmas condições (TANGHE et al., 2002; AUCLAIR et al., 2013). Da mesma forma, na ausência de um oócito, as células do cumulus não são metabolicamente competentes o que altera ou até mesmo impede a execução de vários processos metabólicos, como a expansão do cumulus (RICHANI et al., 2021).


Um dos papéis preponderantes das junções comunicantes é a manutenção do bloqueio meiótico oocitário. Nos mamíferos os oócitos permanecem bloqueados como vesícula germinativa no estágio diplóteno da prófase I da meiose até o pico pré-ovulatório do hormônio luteinizante (LH) (SIRARD et al., 1989; RICHARDS et al., 2002; DEL COLLADO et al., 2018). Através da sinalização regulatória das junções comunicantes ocorre a passagem de monofosfato de adenosina cíclico (AMPc) e o monofosfato de guanosina cíclico (GMPc) para o oócito a partir das células do cumulus. Esses nucleotídeos cíclicos desempenham um papel crítico na regulação meiótica oocitária pois altos níveis intracelulares de AMPc no oócito promovem o bloqueio meiótico (CONTI et al., 2012), portanto a degradação do AMPc presente no oócito é essencial para a retomada da meiose. Essa degradação é feita pela ação da enzima AMPc-fosfodiesterase (PDE), contudo, a transferência de GMPc das células do cumulus para oócito através das junções comunicantes inibe a ativação da PDE, mantendo os

níveis de AMPc intraocitários altos e, conseqüentemente, o bloqueio meiótico (SELA-ABRAMOVICH et al., 2006).

O pico de LH liberado pela adenohipófise determina redução na produção GMPc e, com a retomada da meiose e expansão do cumulus, concomitantemente ocorre o desacoplamento das junções comunicantes. Em conjunto esses eventos determinam a diminuição do fluxo de GMPc para o oócito, permitindo a rápida degradação do AMPc pela PDE e a consequente retomada da meiose in vivo (SUN; MIAO; SCHATTEN, 2009). In vitro a remoção do oócito do folículo ovariano desencadeia espontaneamente a retomada da meiose (LONERGAN et al., 2003).

Apesar de depender diretamente dos íons, metabólitos, aminoácidos e pequenas moléculas reguladoras que recebe das células do cumulus, o oócito regula vários processos nas células somáticas que o rodeiam por meio da secreção dos chamados fatores secretados pelo oócito, dentre os quais o fator de diferenciação de crescimento 9 (GDF9), a proteína morfogenética óssea 15 (BMP15) e alguns membros da família dos fatores de crescimento fibroblásticos (FGF) são os mais estudados, embora não sejam os únicos conhecidos (CAIXETA et al., 2013a; RUSSELL et al., 2016). Por meio desses fatores parácrinos, o oócito é capaz de controlar: a diferenciação fenotípica das células do cumulus em relação as células da granulosa mural, a proliferação das células da granulosa e do cumulus (RUSSELL et al., 2016), a glicólise nas células do cumulus (SUGIURA et al., 2008), a captação de glicose pelo CCO (CAIXETA et al., 2013a), a aquisição da capacidade de sinalização da família EGF nas células do cumulus (SUGIMURA; RICHANI; GILCHRIST, 2018), e a mucificação e expansão das células do cumulus (CAIXETA et al., 2013a; MACHADO et al., 2015). Um esquema gráfico resumido de todo esse processo está representado na Figura 1.

Figura 1 – Representação gráfica da comunicação bidirecional entre o oócito e as células do cumulus



Fonte: Adaptado de (RUSSELL et al., 2016)

2.3 A EXPANSÃO DAS CÉLULAS DO CUMULUS

A ovulação é um processo complexo que determina alterações estruturais, endócrinas e bioquímicas no folículo pré-ovulatório a fim de liberar um gameta feminino apto para ser fecundado e que dê suporte ao futuro embrião durante o período pré-implantacional (SHIMADA et al., 2006). Assim como na retomada da meiose pelo oócito, o fator desencadeante da ovulação é o pico de LH. Esse hormônio se liga a receptores de LH (LHR) nas células da granulosa uma vez que nem o oócito, nem as células do cumulus expressam LHR, e estimula a transcrição de genes relacionados com a ovulação em si, com a expansão do cumulus, com a maturação nuclear e com a esteroidogênese (RICHARDS et al., 2002).

O pico pré-ovulatório de LH também induz a expansão in vivo dos CCOs. Durante a expansão, as células do cumulus secretam uma matriz rica em ácido hialurônico e sintetizam as proteínas estabilizadoras dessa matriz. A hidratação desse ácido promove o aumento no espaço entre as células do cumulus e mucificação da matriz que as envolve (NAGYOVA et al., 2008; YAMASHITA; HISHINUMA; SHIMADA, 2009).

Sob efeito do LH, as células da granulosa produzem fatores de crescimento semelhantes ao EGF (EGF-Like), especialmente epiregulina (EREG), anfiregulina (AREG) e betacelulina (SHIMADA et al., 2006). Esses EGF-Like, particularmente EREG e AREG nos bovinos (PORTELA et al., 2011), atuam de forma autócrina nas células da granulosa mural e de forma parácrina nas células do cumulus. Da mesma forma que o LH nas células da granulosa, o hormônio folículo estimulante (FSH) induz a expressão dos EGF-like nas células do cumulus (CAIXETA et al., 2013b). Entretanto, os EGF-Like a princípio são sintetizados como precursores ancorados à membrana e de forma subsequente são liberados da célula por proteólise mediada por ADAM17 (BEN-AMI et al., 2006). Uma vez liberados, os EGF-Like se ligam a receptores de EGF (EGFR) expressos nas células da granulosa e do cumulus onde estimulam a sua própria síntese, a produção de prostaglandina sintetase 2 (PTGS2) e prostaglandina E2 (PGE2), além de induzir a expressão de genes envolvidos com a esteroidogênese e daqueles envolvidos com a expansão do cumulus. A PGE2 também participa do processo de ovulação por induzir a síntese de EGF-like de maneira semelhante ao LH (SHIMADA et al., 2006).

Recentemente foi demonstrado que a atividade transcricional de *Yes-associated protein* (YAP), um efetor da via Hippo, regula a cascata de sinalização do EGF em células da granulosa mural e do cumulus em bovinos, interferindo diretamente nos processos de

expansão do cumulus e ovulação induzidos pelo LH (DOS SANTOS et al., 2022; KOCH et al., 2022). Dessa forma e em conjunto, os EGF-like, a YAP, a PTGS2 e a PGE2 potencializam a produção da proteína indutora do fator de necrose tumoral 6 (TNFAIP6), da hialurona sintetase 2 (HAS2) e da pentraxina 3 (PTX3) as quais, por sua vez, são responsáveis pela expansão do cumulus (CONTI et al., 2006; SHIMADA et al., 2006; DOS SANTOS et al., 2022; KOCH et al., 2022).

HAS2 promove a síntese do ácido hialurônico, principal componente da matriz extracelular do cumulus (VELHO et al., 2013). TNFAIP6 é necessário para a estabilização da matriz do cumulus expandido e sua expressão é induzida nas células murais da granulosa e nas células do cumulus (NAGYOVA et al., 2008). Essa proteína é indispensável para a fertilidade feminina visto que camundongos deficientes em TNFAIP6 não são capazes de estruturar a matriz extracelular do oócito e, por isso, são estéreis (FÜLÖP et al., 2003). A PTX3 também é essencial para a estabilidade da matriz do cumulus. Sua afinidade pela TNFAIP6 e a interação entre elas parece ser determinante para a estruturação da matriz do cumulus e fertilidade da fêmea (SALUSTRI et al., 2004).

A expressão de mRNA de *HAS2*, *PTGS2*, *TNFAIP6* e *PTX3* foi positivamente associada com aumento na qualidade embrionária e por isso esses genes foram sugeridos como marcadores de competência oocitária nas células do cumulus (NAGYOVA et al., 2008; TESFAYE et al., 2009), assim como a expressão de mRNA dos genes envolvidos na cascata periovulatória (EGF-Likes, *PTGS2*, *YAP*) em células da granulosa é crítica para ruptura da parede do folículo e a liberação do CCO (PORTELA et al., 2011; DOS SANTOS et al., 2022). Assim, a alteração no padrão de expressão de qualquer desses genes sugere que parte das funções ovarianas pode estar comprometida.

2.4 VIAS DE SINALIZAÇÃO E REPARO DOS DANOS DE DNA

A replicação e transmissão precisas do material genético são essenciais para a homeostase celular e viabilidade dos organismos. No entanto, as células estão continuamente expostas a fatores ambientais e agentes endógenos deletérios que prejudicam a integridade do DNA e ameaçam a integridade genômica (PANIER; BOULTON, 2014).

A hidrólise que leva à depuração espontânea do DNA, as espécies reativas de oxigênio (ROS) que induzem oxidação de bases e quebras de DNA, os defeitos de replicação, e o colapso da forquilha de replicação que pode resultar em quebras de fita do DNA são exemplos de processos endógenos prejudiciais ao ácido nucleico (BRANZEI; FOIANI, 2008).

Danos ambientais/exógenos ao DNA podem ser produzidos por fontes físicas ou químicas. A radiação ionizante, tanto de origem cósmica quanto de origem hospitalar (radioterapia ou raios-X) e a luz ultravioleta solar são agentes genotóxicos físicos que podem induzir a oxidação de bases de DNA e gerar quebras de DNA de fita simples e de fita dupla. Os agentes usados na quimioterapia do câncer, os inibidores da topoisomerase que induzem a formação quebras das fitas do DNA, o tabagismo que causa uma grande variedade de adutos e danos oxidativos no pulmão e em outros tecidos, e os APs que também formam adutos com DNA e proteínas, são exemplos de danos de origem química que acometem o DNA (FU et al., 2004; BRANZEI; FOIANI, 2008; LORD; ASHWORTH, 2012). A maturação in vitro (IVM) de CCOs bovinos, especialmente em meio não suplementado com soro fetal bovino (FCS), assim como a vitrificação de embriões murinos determinaram aumento na expressão de genes relacionados com o reparo de DNA (IKEDA; IMAI; YAMADA, 2003; CHANG et al., 2019). Da mesma forma, a exposição a luz ultravioleta induziu a quebra da fita dupla do DNA (DSB) de embriões bovinos produzidos in vitro (BARRETA et al., 2012).

Para proteger apropriadamente o genoma, todos os tipos de alterações estruturais do DNA precisam ser detectadas e corretamente reparadas. A resposta ao dano de DNA (DDR) é executada por uma série de vias de transdução de sinal entrelaçadas que detectam o dano no DNA e, após a ativação dos pontos de verificação do ciclo celular (*cell-cycle checkpoints*), determinam dois desfechos principais: o próprio mecanismo de reparo e/ou apoptose (MAGNUSON; BEDI; LJUNGMAN, 2016). Por meio de uma cascata de sinalização firmemente coordenada, os pontos de verificação do ciclo celular coordenam a parada do ciclo celular e o recrutamento de proteínas de reparo que modificam quimicamente o DNA a fim de proteger o genoma. Entre as proteínas de reparo estão nucleases, helicases, polimerases, topoisomerases, recombinases, ligases, glicosilases, demetilases, quinases e fosfatases, as quais devem ser precisamente reguladas já que cada uma, por si só, tem potencial de alterar a integridade do DNA se acionada no momento inapropriado ou no segmento incorreto do ácido nucleico. Uma vez que o DNA tenha sido reparado o ciclo celular é retomado, caso contrário, a DDR encaminha para a senescência ou apoptose celular (CICCIA; ELLEDGE, 2010).

Como outras vias de transdução de sinal, a DDR consiste em sensores de sinal, transdutores e efetores. Os sensores desta via são proteínas que reconhecem estruturas aberrantes, induzidas por danos e/ou estresse de replicação do DNA, e ativam os transdutores. Os transdutores são representados por uma cascata de proteínas quinases e proteínas

mediadoras que facilitam eventos de fosforilação dentro da rede de reparo ao dano no DNA. Os efetores são substratos das quinases transdutoras que participam de um amplo espectro de processos celulares importantes para a estabilidade genômica, como replicação de DNA, reparo de DNA e controle do ciclo celular (MARÉCHAL; ZOU, 2013).

A DDR é mediada principalmente por proteínas (transdutores) da família das proteínas quinases semelhantes a fosfatidilinositol-3 quinase (PIKK) e pelos membros da família poli(ADP-ribose) polimerase (PARP). Nos mamíferos, três quinases da família PIKK atuam como transdutoras: *ataxia telangiectasia mutated* (ATM), *ATM and RAD3 related* (ATR) e subunidade catalítica da proteína-quinase DNA-dependente (DNAPK-cs). A família PARP é composta por 16 membros, porém apenas PARP1 e PARP2 estão envolvidos na DDR (CICCIA; ELLEDGE, 2010).

Em resposta ao dano do DNA, centenas de proteínas podem ser fosforiladas por ATM ou ATR, enquanto a DNAPK parece regular um número menor de substratos. ATM e DNAPK são ativados por agentes que determinam DSBs. Por sua vez, ATR responde a um amplo espectro de danos ao DNA, incluindo DSBs e uma gama de lesões de DNA que interferem na replicação. PARP1 e PARP2 são ativados tanto por quebras de fita simples no DNA (SSB)s quanto por DSBs e catalisam a adição de cadeias de poli(ADP-ribose) em proteínas para recrutar fatores para a DDR (MARÉCHAL; ZOU, 2013).

A proteína de replicação A (RPA) tem alta afinidade por regiões de DNA de fita simples e, quando ocorrem SSBs, sua ligação a esses substratos parece ser o sinal para o recrutamento das quinases transdutoras. Por sua vez, ATM é recrutada poucos minutos após a ocorrência de DSB e fosforila várias proteínas que são necessárias para sinalizar danos e reparos no DNA. A histona H2AX é um alvo proximal de atividade nessa cascata de sinalização e, após ser fosforilada por ATM, DNAPK ou ATR no local do dano, age no recrutamento de proteínas que iniciam o processo de remodelação da cromatina. ATM e ATR também atuam reduzindo a atividade da quinase dependente de ciclina, pela ativação da transcrição de p53. Isso causa um atraso na progressão do ciclo celular, aumentando o tempo disponível para que ocorra o reparo das lesões do DNA antes da replicação ou divisão celular (CICCIA; ELLEDGE, 2010; CECCALDI; RONDINELLI; D'ANDREA, 2016).

A seleção de qual sistema de reparo será ativado depende do tipo de lesão no DNA e da fase do ciclo celular que o dano foi detectado (BRANZEI; FOIANI, 2008). A via de reparo por excisão de nucleotídeos (*nucleotide excision repair*) atua no reparo a danos que causam a distorção da hélice do DNA e que interferem no pareamento das bases. Dessa forma, quando

esses erros não são reparados ocorre a interrupção da transcrição e a replicação normais. A via do reparo por excisão de bases (*base excision repair*) é responsável por corrigir pequenas alterações nas bases nitrogenadas. Essas alterações geralmente não impedem a transcrição ou a replicação apesar de muitas vezes gerarem erros de codificação. A via do reparo a erros de pareamento (*mismatch repair*) é responsável pela correção de erros de pareamento das bases nitrogenadas e pequenas inserções ou deleções que ocorrem durante a replicação celular. As lesões que causam quebra física de uma ou de ambas as fitas de DNA são alvo de dois tipos de sistemas de reparo. O primeiro independe de homologia entre as fitas do DNA e é representado pela via da união terminal não homóloga (NHEJ - *non-homologous end-joining*); o segundo é representado por três vias dependentes de homologia: recombinação homóloga (HR - *homologous recombination*), união terminal mediada por micro-homologia (MMEJ - *microhomology-mediated end joining*) e anelamento de fitas simples (SSA - *single-strand annealing*) (CICCIA; ELLEDGE, 2010; LORD; ASHWORTH, 2012).

2.4.1 Reparos de quebras na fita dupla do DNA

Dos muitos tipos de lesões de DNA, as DSBs são consideradas as mais prejudiciais já que apenas uma DSB, caso não seja devidamente reparada, é suficiente para desencadear a parada permanente do ciclo e a morte celular. Além disso, as DSBs são potentes indutores de rearranjos cromossômicos, tais como deleções, translocações e amplificações. Esses rearranjos do genoma podem resultar na ativação de oncogenes e/ou na perda de supressores de tumor, o que, por sua vez, alimenta a transformação carcinogênica (PANIER; BOULTON, 2014; SUN et al., 2018).


As DSB são caracterizadas pela quebra simultânea de ligações fosfodiéster na mesma região ou em regiões próximas de ambas as fitas de DNA. Como resultado dessa dupla ruptura, a integridade sequencial da informação genética é fisicamente desconectada, assim como a perda da referência precisa que é utilizada em muitas vias onde apenas uma das fitas está comprometida. Quando esse tipo de lesão é comparado com os demais, fica claro seu nível de complexidade e potencial de dano (JACKSON; BARTEK, 2009; LIEBER, 2010).

Frente a grande diversidade de fatores determinantes de DSBs, algumas vias de reparo especializadas evoluíram para manter a integridade do genoma. As células respondem as DSBs montando uma rede de sinalização complexa que coordena as reações de reparo do DNA com a ativação do ponto de verificação de danos no DNA e a reorganização da cromatina. A maioria desses processos de reparo envolve uma nuclease para remover o DNA

danificado, uma DNA polimerase para preencher o novo DNA e uma ligase para restaurar a integridade de cada fita do duplex (LIEBER et al., 2010). Além disso, pelo menos quatro sensores parcialmente independentes podem detectar DSBs: PARP, Ku70/Ku80, complexo Mre11-Rad50-Nbs1 (MRN) e RPA (CICCIA; ELLEDGE, 2010; CECCALDI; RONDINELLI; D'ANDREA, 2016).

Embora os mecanismos celulares que determinam a escolha da via para o reparo das DSBs ainda não estejam totalmente elucidados, parece que a extensão do processamento das pontas de DNA desempenha papel fundamental nessa escolha (PANIER; BOULTON, 2014). NHEJ não requer ressecção das pontas das fitas de DNA enquanto as outras três vias dependem dessa ressecção, que é mais restritiva (5 – 25 nucleotídeos) para MMEJ e mais extensiva para HR e SSA (CICCIA; ELLEDGE, 2010; CECCALDI; RONDINELLI; D'ANDREA, 2016). A Figura 2 representa estas quatro abordagens para o reparo das DSB.

Figura 2 – Quatro abordagens para o reparo das quebras na fita dupla do DNA



Fonte: Adaptado de (CECCALDI; RONDINELLI; D'ANDREA, 2016)

Por serem as duas principais vias de reparo de DSB (BRANZEI; FOIANI, 2008; JACKSON; BARTEK, 2009; LORD; ASHWORTH, 2012; PANIER; BOULTON, 2014; BOHRER et al., 2018), NHEJ e HR serão descritas a seguir.

2.4.1.1 União terminal não homóloga (NHEJ)

A NHEJ representa o mecanismo mais simples e rápido para reparar DSBs, portanto, é a via de reparo predominante na maioria das células de mamíferos, embora possa ocasionalmente levar à perda de informação genética (VÍTOR et al., 2020). NHEJ promove a ligação entre as pontas das fitas de DNA rompidas independente de homologia e/ou grande ressecção, portanto, essa via está mais propensa a erros que a HR. Pode ser acionada durante qualquer uma das fases do ciclo celular (LIEBER, 2010; LORD; ASHWORTH, 2012) mas a alta compactação da cromatina e a ausência de cromátides irmãs disponíveis são importantes fatores que tornam a NHEJ a via de reparo predominante durante a fase G1 (WARD et al., 2004; BRANZEI; FOIANI, 2008; FENG et al., 2015; VÍTOR et al., 2020), contudo, para Ceccaldi, Rondinelli e D'Andrea (2016) essa é a via dominante nas fases G0/G1 e G2 do ciclo celular.

O ponto de partida do processo NHEJ se dá com o reconhecimento da DSB e ligação das extremidades do DNA de fita dupla pelo heterodímero KU, formado pelas proteínas KU70 e KU80 (produtos dos genes *XRCC6* e *XRCC5*, respectivamente), o que ocorre de maneira extraordinariamente rápida e eficiente devido à sua abundância e forte atração por esse tipo de substrato de DNA. O heterodímero KU ligado ao DNA, por sua vez, recruta DNAPK-cs para formar a holoenzima DNA-PK, de modo que as duas moléculas de DNAPK-cs ligadas a lados opostos do DSB possam interagir entre si, contribuindo para a sinapse das extremidades quebradas do DNA e impedindo a ressecção das extremidades das fitas rompidas. O complexo DNA-PK é o principal regulador do processo NHEJ, coordenando o recrutamento de fatores acessórios, como a proteína 4 de complementação cruzada de reparo de raios X (*XRCC4*) e a DNA ligase IV (*LIG4*), que contribuem para o adequado pareamento das extremidades DSB e realizam a ligação final das fitas de DNA (CICCIA; ELLEDGE, 2010; SERRANO-BENÍTEZ; CORTÉS-LEDESMA; RUIZ, 2020).

Após a ligação da DSB, a auto fosforilação de DNAPK-cs resulta na desestabilização de sua interação com as extremidades do DNA, possibilitando acesso a enzimas de processamento terminal, como ARTEMIS. Essa auto fosforilação da DNAPK-cs, em seu cluster PQR, previne o processamento terminal excessivo e tem efeito inibitório sobre a HR pois impede a ressecção excessiva da dupla fita (CICCIA; ELLEDGE, 2010).

2.4.1.2 Recombinação homóloga (HR)

A HR requer uma sequência de DNA homóloga para servir de molde para o reparo das DSB e envolve extensivo processamento do DNA. Como esperado, a HR é extremamente precisa, pois leva ao reparo do locus danificado usando sequências de DNA homólogas não danificadas às extremidades quebradas. Há fortes indícios de que a ressecção das extremidades das fitas de DNA no sentido 5' – 3', promovendo a extensão da extremidade 3'ssDNA (fita simples de DNA), é um passo crítico e determinante para escolha da HR como via de reparo para as DSBs. Esses trechos de 3'ssDNA estendidos são utilizados para busca da região de homologia na cromátide irmã que servirá de molde para o reparo (MARINI et al., 2019; VÍTOR et al., 2020). HR acontece principalmente nas fases S e G2 do ciclo celular, momentos nos quais a replicação do DNA é mais intensa e existe uma cromátide irmã disponível (BRANZEI; FOIANI, 2008; LORD; ASHWORTH, 2012; FENG et al., 2015; VÍTOR et al., 2020).

A via HR pode ser dividida didaticamente em três etapas: pré-sinapse, sinapse e pós-sinapse. Na pré-sinapse, ocorre o reconhecimento das quebras de fita dupla e posterior processamento das pontas do DNA (ressecção), gerando filamentos de ssDNA em cada uma das fitas quebradas. Na fase sináptica ocorre a invasão da fita de DNA a ser reparada em seu molde de reparo na cromátide homóloga. Esta invasão resulta em uma estrutura chamada de D-loop. A seguir inicia-se a fase pós sinapse quando ocorre a síntese da porção danificada e a separação das cromátides homólogas (RENKAWITZ; LADEMANN; JENTSCH, 2014).

A HR é iniciada pelo reconhecimento das DSBs pelo complexo MRE11-RAD50-NBS1 (MRN), que promove a ativação de ATM e a preparação de DNA para o reparo. RAD50 contém domínios ATPase que interagem com MRE11 e se associam com as extremidades rompidas de DNA. MRE11 também possui atividades de endonuclease e exonuclease que atuam nas etapas iniciais de ressecção das fitas danificadas. A terceira subunidade do complexo MRN, NBS1, associa-se com ATM através de sua região C-terminal, que promove o recrutamento de ATM para DSBs, onde ATM é ativado pelo complexo MRN (CICCIA; ELLEDGE, 2010).

A ressecção final do DNA é regulada por ATM através da proteína *CtBP-interacting* (CtIP), que interage com BRCA1 e MRN. Durante este processo, a formação de DNA de fita simples (ssDNA) é continuada pela ação da exonuclease 1 (Exo1). Em S e G2, CtIP associa-se com BRCA1 que facilita sua ligação com locais de dano. Na fase G1 CtIP atua de maneira independente de BRCA1 e, dessa forma, promove a reparação por MMEJ que é mediada pelo

emparelhamento de regiões de microhomologia de ssDNA, seguido por ligação terminal de DNA dependente de Ligase 3 (CICCIA; ELLEDGE, 2010; RENKAWITZ; LADEMANN; JENTSCH, 2014)

Na sequência do processo a RPA recobre a extremidade da ssDNA protegendo-a da formação de estruturas secundárias (GUDMUNDSDOTTIR; ASHWORTH, 2006; MARINI et al., 2019). Além da estabilidade das extremidades do DNA, RPA atua garantindo que haja a correta incorporação de filamentos RAD51 (*DNA repair protein RAD51*), mediada por BRCA2 (*Breast Cancer 2 DNA repair associated*) e RAD52 (*DNA repair protein RAD52*) (CICCIA; ELLEDGE, 2010). Por sua vez, durante a fase sináptica, o filamento RAD51 invade a cadeia de DNA não danificada e pareia com um duplex homólogo, se uma sequência doadora tiver sido encontrada durante o processo de pesquisa de homologia. A troca de fitas de DNA entre o DNA alvo e o filamento RAD51 gera uma estrutura conhecida como loop de deslocamento (D-loop) que contém o novo DNA heteroduplex e a fita deslocada do DNA doador. Por fim a síntese de DNA é iniciada a partir da extremidade 3', a segunda extremidade de DSB alinha-se com o D-loop estendido para formar a dupla junção *Holliday* e as estruturas resultantes são resolvidas. Após a síntese as junções *Holladays* são clivadas e as fitas de DNA são ligadas para produzir duas moléculas intactas de DNA (CICCIA; ELLEDGE, 2010; RENKAWITZ; LADEMANN; JENTSCH, 2014; CECCALDI; RONDINELLI; D'ANDREA, 2016).

A Proteína de Ligação de Proteína Tumoral P53 1 (TP53BP1) é um importante regulador da resposta celular a DSB. Apresenta marcado antagonismo com BRCA1 e, portanto, o equilíbrio entre BRCA1 e TP53BP1 é o modulador para determinação da via de reparo que será utilizada. Como TP53BP1 bloqueia a ressecção de DNA, impedindo que o CtIP e ATM acessem as extremidades do ácido nucleico durante a fase G1 do ciclo celular, sua atuação promove a via NHEJ em detrimento a HR (CECCALDI; RONDINELLI; D'ANDREA, 2016; MARINI et al., 2019). Dessa forma, TP53BP1 bloqueia a via HR e promove a NHEJ. Contudo, essa proteção das extremidades do DNA depende da fosforilação da TP53BP1 por ATM que promove o recrutamento de fatores de interação *Rap1-interacting factor 1* (RIF1) e *PAX-interacting protein 1* (PTIP) (SERRANO-BENÍTEZ; CORTÉS-LEDESMA; RUIZ, 2020). Vale ressaltar que nas fases S/G2 do ciclo celular, BRCA1 promove a desfosforilação de 53BP1 e a liberação de RIF1, determinando a ressecção das extremidades das fitas do DNA e direcionando o reparo para HR (MARINI et al., 2019).

Os principais genes que controlam as vias HR e NHEJ de reparo de DNA são expressos em embriões bovinos (BARRETA et al., 2012) e oócitos e blastocistos humanos (JAROUDI et al., 2009). Embriões bovinos com baixa competência mostraram maior expressão de *TP53BP1* e *RAD52* quando comparados aos embriões com alta competência. Quando DSBs foram induzidas experimentalmente por radiação ultravioleta em blastocistos bovinos, a expressão de *TP53BP1*, *RAD51* e *KU70* diminuiu 72 horas pós inseminação e depois ficou maior que a do grupo não irradiado as 168 horas pós inseminação (BARRETA et al., 2012). Em conjunto esses resultados indicam que o perfil de expressão dos genes envolvidos no reparo das DSBs em embriões com baixa competência está alterado e, portanto, podem atuar como indicadores de viabilidade e saúde dos embriões.

ARTIGO I

ARTIGO A SER SUBMETIDO PARA PUBLICAÇÃO

***Senecio brasiliensis*-DERIVED PYRROLIZIDINE ALKALOIDS DECREASE
OVULATORY CASCADE GENE EXPRESSION AND BOVINE EMBRYOS
PRODUCTION**

André Lucio Fontana Goetten, Carolina dos Santos Amaral, Daniele Missio, Vitor Braga
Rissi, Alceu Mezzalira, Christopher Allan Price, Marcos Henrique Barreta & Valério Marques
Portela

REPRODUCTIVE TOXICOLOGY, 2022

***Senecio brasiliensis*-derived pyrrolizidine alkaloids decrease ovulatory cascade gene expression and bovine embryos production**

André Lucio Fontana Goetten^a, Carolina dos Santos Amaral^b, Daniele Missio^b, Alceu Mezzalira^c, Vitor Braga Rissi^a, Christopher Allan Price^d, Gustavo Zamberlam^d, Marcos Henrique Barreta^a & Valério Marques Portela^{b*}

^a Laboratório de Fisiologia da Reprodução Animal (LAFRA), Centro de Ciências Rurais (CCR), Universidade Federal de Santa Catarina (UFSC) - Curitibanos, SC, Brazil.

^b Laboratório de Biotecnologia e Reprodução Animal (BIOREP), Universidade Federal de Santa Maria (UFSM) - Santa Maria, RS, Brazil.

^c Laboratório de Reprodução Animal Assis Roberto de Bem, Centro de Ciências Agroveterinárias (CAV), Universidade do Estado de Santa Catarina (UDESC) - Lages, SC, Brazil.

^d Centre de recherche en reproduction et fertilité (CRRF), Faculté de Médecine Vétérinaire (FMV), Université de Montréal (UdeM), Canada.

* Corresponding author: Valério M. Portela, Laboratório de Biotecnologia e Reprodução Animal, BioRep, Universidade Federal de Santa Maria, Av. Roraima 1000, 97105-900, Santa Maria, RS, Brazil; e-mail: valerio.portela@ufsm.br Santa Maria, RS.

ABSTRACT

Pyrrolizidine alkaloids (PAs)-containing plants are widely distributed in the world and probably the most common poisonous plants affecting livestock, wildlife, and humans. Livestock poisoning, mainly liver damage, depends on PAs metabolism by cytochrome P450, whose isoforms can occur in other organs, including ovaries. This study aimed to assess whether low PAs concentration have a direct effect in vitro on bovine granulosa cells and embryos. Low concentrations (0.03 or 0.3 μ M) of senecionine or seneciphylline, PAs present in the *Senecio* genus, were added to the culture medium. Senecionine and seneciphylline downregulated mRNA expression of *AREG*, *EREG*, *PTGS2*, and *CTGF* genes in granulosa cells, suggesting that PAs could impair bovine ovulation. Both tested PAs do not impair cumulus expansion. Cleavage and blastocyst rates did not differ from the control group when PAs were added to in vitro maturation medium. Cleavage and blastocyst rates were reduced when senecionine or seneciphylline were added to the medium 18–20 hours post-insemination (hpi). Seneciphylline added to maturation medium downregulated the *TP53BP1* gene in blastocysts but upregulated the *RAD51* gene when added at 18–20 hpi, suggesting that PAs could determine DNA double-strand break and homologous repair pathway activation. Senecionine reduced *GLUT3* and *BCL2* mRNA expression, suggesting that PAs could impair glucose metabolism and reduce antiapoptotic activity in bovine blastocysts. In summary, this study demonstrates that senecionine and seneciphylline have significant toxicological effects on bovine granulosa cells as well as in in vitro produced bovine embryos, suggesting that PAs, even at low concentrations, have the potential to directly impact cattle reproductive physiology.

Keywords: Senecionine, seneciphylline, granulosa cells, ovary, embryo.

3.1 INTRODUCTION

Plant intoxication is still one of the most common causes of death in adult production animals, resulting in significant direct and indirect economic losses in the livestock production system [1–3]. Management failures in pasture or crop areas favor the appearance of invasive plants, many of which have toxic potential for various animal species, including humans [4]. The toxic plants in southern Brazil that generate the most economic losses in livestock are those belonging to the genus *Senecio* spp., which cause 50% of all cattle deaths by phytotoxins [2,4,5]. *Senecio brasiliensis* is one of the most dangerous species in the genus.

It can cause hepatotoxicity and poisoning in horses and cattle, which can lead to their death [6].

The toxic principles present in *Senecio* spp. belongs to the group of pyrrolizidine alkaloids (PAs). PAs described in *S. brasiliensis* include senecionine, seneciphylline, integerrimine, retrorsine, usaramine, and jacobine [7,8], with senecionine and seneciphylline being the most prevalent [9]. There is considerable variation in the content of PAs in different plant species or in different parts of the same plant, however, PAs only become toxic when biotransformed into pyrroles, mainly by hepatic cytochrome P-450 enzymes [10]. Pyrroles cause irreversible damage to hepatocytes by inhibiting mitosis and causing necrosis, resulting in hepatic dysfunction [11]. Megalocytosis is one of the most common findings in liver biopsies of cattle that have been intoxicated with *S. brasiliensis*. It could be caused by inhibition of mitosis [12] or by DNA damage [10]. It is already known that some of the pyrroles produced in the liver can reach other organs through the systemic circulation and cause extrahepatic damage [4,13].

Both the acute and chronic forms of intoxication by *S. brasiliensis* in cattle have been experimentally demonstrated. Single doses of 5 to 10 g/kg resulted in chronic poisoning, but when animals received daily repeated doses of 0.625 to 5 g/kg for 1 to 8 months, they did not show any symptoms for weeks or even months, simulating a cumulative effect of these xenobiotics [14]. During the first 18 hours after receiving 1 mg/kg of the alkaloid orally in a single dose, seneciphylline levels greater than 0.3 μ M were found in the blood and milk of cattle. Blood levels of 0.033 μ M were still present after 54 hours of treatment [15]. Given that the amount of PAs in *S. brasiliensis* ranged from 0.178 to 0.31 percent [16,17], an intake of 0.3–0.5 g/kg BW of aerial parts of the plant would be enough to determine blood levels of seneciphylline equivalent to 0.3 μ M.

The decrease in reproductive rates of seneciosis-affected cows has traditionally been linked to a deterioration in nutritional condition caused by the liver damage. However, there are evidences that PAs act directly on ovarian cells since swine ovaries are capable of expressing and transcribing the enzyme cytochrome P-450 and its isoforms [18], and ovaries of bovines with chronic seneciosis showed lesions in large luteal cells similar to those present in hepatocytes [19]. There are also reports that prenatal exposure to APs can impair physical and behavioral development in rats [20].

When we associated cattle infertility with no apparent cause with evidence of PAs acting on reproductive organs, we hypothesized that low PAs concentrations, insufficient to

cause chronic seneciosis, could act directly on ovarian cells and/or embryos, affecting the periovulatory EGF-like cascade and early embryonic development. The aims of this study were to evaluate whether low concentration of pyrrolizidine alkaloids have a direct effect: (a) on expression of periovulatory Epidermal Growth Factor-Like (EGF-Like) cascade genes in bovine granulosa cells; (b) on bovine cumulus-oocyte cell expansion; and (c) on early embryonic development in cattle.

3.2 MATERIAL AND METHODS

All reagents were purchased from Sigma-Aldrich Co. (Rocklin, CA, USA) unless otherwise stated. Senecionine (cat. # 50351) and seneciphylline (cat. # 73913) were dissolved in ethanol, and the final concentration of ethanol never exceeded 0.01% in the media.

3.2.1 Granulosa cell culture

The granulosa cell (GC) culture system employed herein was as a previously described [21,22] in which abundance of mRNA encoding the EGF-like factors and other critical preovulatory genes is acutely upregulated by luteinizing hormone (LH). Bovine ovaries containing large follicles were collected from adult cows, a part of the estrous cycle stage, at a local abattoir and were transported to the laboratory in saline solution (0.9% NaCl; 25 °C) containing penicillin (100 IU/mL), and streptomycin sulfate (50 µg/mL). For each experimental replicate, ten ovaries that each contained a single large follicle (≥ 10 mm diameter) were selected and GC were collected from each large follicle by aspiration, pooled in a single tube and were washed twice by centrifugation at $219\times g$ for 20 min each. Cell viability was estimated with 0.4% Trypan Blue Stain. Cells were then seeded into 24-well tissue culture plates (Sarstedt, St-Leonard, QC, Canada) at a density of 1×10^6 viable cells per well in 1 mL DMEM-F12 supplemented with sodium bicarbonate (10 mM), sodium selenite (4 ng/mL), bovine serum albumin (BSA; 1 mg/mL), penicillin (100 IU/mL), streptomycin (100 µg/mL), transferrin (2.5 µg/mL), non-essential amino acid mix (1.1 mM), androstenedione (10^{-7} M), follicle-stimulating hormone (FSH; 1 ng/mL), insulin (10 ng/mL) and 2% fetal calf serum (FCS). Cultures were maintained at 38.5 °C in 5% CO₂ for 24 h. Medium was then replaced with 0.9 mL serum-free DMEM-F12 with antibiotics for 18 h, at which point LH (100 ng/mL) and the experimental treatments were added in 100 µl DMEM-F12 (with antibiotics).

3.2.2 Cumulus-oocyte complexes recovery and in vitro maturation (IVM)

Bovine ovaries were collected from adult cows, a part of estrous cycle stage, at a local abattoir and were transported to the laboratory in saline solution (0.9% NaCl; 25 °C) containing penicillin (100 IU/mL), and streptomycin sulfate (50 µg/mL). Cumulus oocyte complexes (COCs) from 3 to 8 mm diameter follicles were aspirated with a vacuum pump (vacuum rate of 15 mL of water/minute) and pooled in a 15 mL conical tube. After sedimentation, COCs were recovered and selected under a stereomicroscope as previously described [23]. Grade 1 or 2 COCs were randomly transferred to well plates containing basic maturation medium composed by TCM199 containing Earle's salts and L-glutamine supplemented with 25 mM HEPES, 0.2 mM pyruvic acid, 2.2 mg/mL sodium bicarbonate, 5.0 µg/mL LH (Lutropin-V, Bioniche Animal Health Canada Inc., Belleville, ON, Canada), 0.5 µg/mL FSH (Folltropin-V, Bioniche Animal Health Canada Inc., Belleville, ON, Canada), 100 IU/mL penicillin, 50 µg/mL streptomycin sulfate and 10% FCS. Cultures were maintained in an incubator at 38.5 °C in 5% CO₂. The experimental treatments, number of replicates, and number of COCs/replicate varied according to the objective of each experiment.

3.2.3 In vitro fertilization (IVF) and in vitro embryo culture (IVC)

After IVM, bovine oocytes were fertilized in vitro with frozen-thawed semen and fractionating on discontinuous Percoll (GE Healthcare, São Paulo, SP, Brazil) gradient. Sperm was diluted and added to the COCs plate with final concentration adjusted to 2×10^6 sperm/mL in Fert-TALP medium containing 20 mM penicillinamine, 10mM hypotaurine, 1mM epinephrine [24], 100 IU/mL penicillin, and 50 µg/mL streptomycin sulfate. Fertilization was carried out by coculture of sperm and oocytes for 18–20 hours in four-well plates (Nunc, Roskilde, Denmark) in the same atmospheric conditions used for maturation. IVF day was considered as Day 0 of embryo production.

After IVF, presumptive bovine zygotes were denuded by 2-min vortexing, and then cultured in groups of 35 – 45 in a culture chamber (CBS Scientific, Del Mar, CA) at 38.5°C and saturated humidity atmosphere of 5% CO₂, 5% O₂, and 90% N₂ in 400 µL of synthetic oviduct fluid medium (SOFaaci) [25] containing 0.2mM pyruvic acid, 5% of FCS, 100 IU/mL penicillin and 50 µg/mL streptomycin sulfate in four-well plates. Cleavage rates were evaluated 48 hours post-insemination (hpi), and blastocyst rates were assessed 168 hpi.

Blastocysts evaluated 168 hpi were rinsed three times in phosphate-buffered saline (PBS) and stored at -80°C for subsequent RNA extraction.

3.2.4 RNA extraction, reverse transcription, and real-time polymerase chain reaction

Total RNA was then extracted using PureLink™ RNA Mini Kit (Thermo Fisher Scientific, Waltham, MA, EUA) according to the manufacturer's instructions and was quantified at 260 nm wavelength using a spectrophotometer (NanoDrop1000, Thermo Scientific, Wilmington, DE, USA). Total RNA was treated with 0.1 U DNase Amplification Grade (Thermo Fisher Scientific) for 15 min at 27°C to neutralize any DNA molecules. DNase was inactivated with 1 μl ethylenediaminetetraacetic acid for 10 min at 65°C . Total RNA (200 ng) was reverse transcribed using the iScript™ cDNA Synthesis Kit (Bio-Rad, Des Plaines, IL, USA) at 25°C for 5 min and 42°C for 30 min. The reaction was ended by incubation at 85°C for 5 min.

Real-time qPCR was performed using CFX384™ Real-Time System (Bio-Rad Laboratories, Hercules, CA, USA) using 2 μl of complementary DNA (cDNA) and 8 μl of MIX containing forward and reverse bovine specific primers (Table 1), nucleases free water, and GoTaq® Master Mix (Promega Corporation, Madison). Amplification was performed with an initial denaturation at 95°C for 5 min followed by 40 cycles of denaturation at 95°C for 15 s and annealing/extension at 60°C for 30 s. Samples were run in duplicate. To optimize the RT-qPCR assays, serial dilutions of cDNA templates were used to generate a standard curve, and efficiency between 90% and 110% and coefficient of determination (R^2) higher than 0.98 were considered. To select the most stable housekeeping gene H2A histone family, member Z (*H2AFZ*), cyclophilin B (*PPIB*), TATA-box binding protein (*TBP*), glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*), and ribosomal protein S18 (*RPS18*) amplification profiles were compared using the geNorm software [26]. The results indicated that *GAPDH* was the most stable endogenous control for both blastocysts and granulosa cells. The expression of each target gene, relative to the expression of the reference genes, was calculated using the Pfaffl method with correction for amplification efficiency and normalized to a calibrator sample [27].

Table 1 – Details of primers used for gene expression analysis RT-qPCR

Gene Symbol	Primer sequence (5' → 3')	Accession Number
<i>H2AFZ</i>	F: GAGGAGCTGAACAAGCTGTTG R: TTGTGGTGGCTCTCAGTCTTC	XM_002686087.4

Gene Symbol	Primer sequence (5' → 3')	Accession Number
<i>PPIB</i>	F: GGTCATCGGTCTCTTTGGAA R: TCCTTGATCACACGATGGAA	NM_174152.2
<i>TBP</i>	F: CAGAGAGCTCCGGGATCGT R: CACCATCTTCCCAGAACTGAATAT	NM_001075742.1
<i>GAPDH</i>	F: GATTGTCAGCAATGCCTCCT R: GGTCATAAGTCCCTCCACGA	NM_001034034.2
<i>RPS18</i>	F: CCTTCCGCGAGGATCCATTG R: CGCTCCCAAGATCCAACTAC	NM_001033614.2
<i>AREG</i>	F: CTTTCGTCTCTGCCATGACCTT R: CGTTCTTCAGCGACACCTTCA	NM_001099092.1
<i>EREG</i>	F: CTGCACAGCATTAGTTCAAACCTGA R: TGTCCATGCAAACAGTAGCCATT	XM_010806226.3
<i>PTGS2</i>	F: TGCTGAGTTTAACACGCTCTACCA R: TGAGACCATGTTCCAGTAAGACAGA	NM_174445.2
<i>CTGF</i>	F: AGCTGAGCGAGTTGTGTACC R: TCCGAAAATGTAGGGGGGCAC	NM_174030.2
<i>TNFAIP6</i>	F: GCAAAGGAGTGTGGTGGTGTGTTT R: ACTGAGGTGAATGCGCTGACCATA	NM_001007813.2
<i>GADD45B</i>	F: TACGAGTCGGCCAAGCTGAT R: GTCCTCCTCTTCCTCGTCGAT	NM_001040604.1
<i>TP53BP1</i>	F: ATCAGACCAACAGCAGAATTTCC R: CACCACGTCAAACACCCCTAA	NM_001206397.2
<i>XRCC6</i>	F: AATTGACTCCTTTTGACATGAGCAT R: CCATAGAACACCACTGCCAAGA	NM_001192246.1
<i>RAD51</i>	F: ATGCACCGAAGAAGGAGCTAAT R: GATCGCCTTTGGTGGAACTC	NM_001046179.2
<i>BAX</i>	F: GACATTGGACTTCCTTCGAGA R: AGCACTCCAGCCACAAAGAT	NM_173894.1

Gene Symbol	Primer sequence (5' → 3')	Accession Number
<i>BCL2</i>	F: GTGGATGACCGAGTACCTGAAC R: AGACAGCCAGGAGAAATCAAAC	NM_001166486
<i>GLUT1</i>	F: CCTTCACTGTCGTGTCGCTA R: GCCACAATGCTCAGGTAGGA	NM_174602.2
<i>GLUT3</i>	F: GCCGCCGATAGAGGACATTT R: ATGGCGAAGATCAGAGGTGC	NM_174603.3

3.3 STATISTICAL ANALYSIS

The effects of PAs on gene expression in GCs were assessed using a two-way analysis of variance (ANOVA). One-way ANOVA was used to test the effects of PAs on gene expression in blastocysts. The effects of PAs on cumulus expansion were investigated using a two-way repeated measures ANOVA analysis. Differences were significant when $P < 0.05$. When a significant group effect was detected, comparisons were performed using the least-square means (LSMeans) method (R Package “emmeans” version 1.6.3) with Tukey-Kramer HSD test for multiple comparisons adjustment. The Shapiro-Wilk test was used to check the assumption of normality, and the Bartlett test was used to test the homogeneity of variances. The Box-Cox power transformation was used to transform data that did not follow a normal distribution or showed heteroscedasticity. All data are presented as means $\pm$ standard error of mean (SEM) and the level of significance was set at $P < 0.05$. Cleavage and blastocyst rates were examined by a binomial generalized linear model, with log link. Risk ratios (RRs) and 95% confidence intervals (CIs) were used to present the result. All statistical analyzes were performed using the R 4.1.1 software (R Core Team, 2021, Vienna, Austria).

3.3.1 Experimental design

3.3.1.1 Experiment 1 – Effects of PAs on the periovulatory EGF-Like cascade genes in granulosa cells

To assess the effects of Senecionine and Seneciphylline in graded concentrations (0, 0.03, and 0.3 μ M) on amphiregulin (AREG), epiregulin (EREG), prostaglandin-endoperoxide synthase 2 (PTGS2), and connective tissue growth factor (CTGF) mRNA expression, GCs were cultured as described in the section 3.2.1 and randomly distributed into the following

groups: Negative Control (Neg_Ctrl; without PAs and LH), Positive Control (Pos_Ctrl; without PAs), Senecionine 0.03 μ M (Cion_0.03), Senecionine 0.3 μ M (Cion_0.3), Seneciphylline 0.03 μ M (Phyl_0.03), and Seneciphylline 0.3 μ M (Phyl_0.3). At 6, 12, and 24 h post-treatment culture medium was removed and GCs were collected for total RNA extraction and RT-qPCR analyses. The main effects of group (PAs graded concentration), time (6, 12, and 24 h post-treatment), and their interaction were tested. This experiment was performed in three independent replicates, with each replicate using ovaries collected at different days.

3.3.1.2 Experiment 2 - Effect of PAs on cumulus-oocyte complexes expansion

To study the effect of senecionine or seneciphylline inclusion in graded concentrations (0, 0.03, and 0.3 μ M) during in vitro maturation on COCs expansion, COCs (n=8/group/replicate) were randomly distributed into the following groups: Negative Control (Neg_Ctrl; without PAs, FSH, and LH), Positive Control (Pos_Ctr; without PAs), Senecionine 0.03 μ M (Cion_0.03), Senecionine 0.3 μ M (Cion_0.3), Seneciphylline 0.03 μ M (Phyl_0.03), and Seneciphylline 0.3 μ M (Phyl_0.3). COCs were individually (1 COC/well) matured in 50 μ L of maturation medium in 96-well plates (Sarstedt, St-Leonard, QC, Canada). COCs from each treatment group were photographed using an inverted microscope (Nikon Eclipse TS-100; Nikon Instruments Inc., Melville, NY). Images of each COC were captured through Micrometrics SE Premium software (version 4.5.1, Unitron, Commack, NY, USA) at 0, 6, 12, and 24 h of maturation. Total surface area of each COC expressed in pixels was measured with ImageJ software (version 1.50, National Institutes of Health, Bethesda, MD, USA). The relative expansion (RE) of each COC at a given time point was calculated by dividing the area of the COC at that time point by the area at time 0. The main effects of group (PAs graded concentration), time (6, 12, and 24 h of IVM), and their interaction were tested. This experiment was performed in three independent replicates, with each replicate using ovaries collected at different days.

3.3.1.3 Experiment 3 - Effects of PAs added to the IVM medium on bovine embryo production and quality.

The experimental design used in IVM was similar to the experiment 2 with small modifications. COCs (n = 40 – 45/group/replicate) were matured in four-well plates (Nunc) containing 400 μ L of maturation medium with graded concentrations (0, 0.03 μ M, and 0.3 μ M) of both PAs per well. The Neg_Ctrl group was not used.

Following IVM, oocytes were fertilized and presumptive zygotes were cultured in media free of PAs, as described in section 3.2.3. To assess embryo production, cleavage rates were evaluated 48 hpi, and blastocyst rates were assessed 168 hpi. Blastocysts evaluated 168 hpi were rinsed three times in phosphate-buffered saline (PBS) and collected for total RNA extraction and RT-qPCR analyses. To assess embryo quality, mRNA expression of glucose-1-transporter (*GLUT1*), glucose-3-transporter (*GLUT3*), tumor protein p53 binding protein 1 (*TP53BP1*), RAD51 recombinase (*RAD51*), X-ray repair cross complementing 6 (*XRCC6*), B-cell lymphoma 2 (*BCL2*), and BCL2 associated X (*BAX*) were determined. This experiment was performed in three independent replicates, with each replicate using ovaries collected at different days.

3.3.1.4 Experiment 4 - Effects of PAs added to the IVC medium on bovine embryo production and quality.

The experimental design used in this experiment was similar to the experiment 3 with small modifications. No PAs were added to the medium during maturation or fertilization. Only after denudation (18-20 hpi), were the presumptive zygotes cultured in medium containing PAs at the same concentrations used in previous experiments. Embryo production and quality were assessed in the same manner as in experiment 3. This experiment was performed in five independent replicates, with each replicate using ovaries collected at different days.

3.4 RESULTS

3.4.1 mRNA relative expression of periovulatory EGF-Like cascade genes in bovine granulosa cells exposed to PAs

The transcription levels of all genes tested in the positive control group increased in a time-dependent manner following LH exposure, as expected for this model, whereas in the absence of LH stimulation (negative control group), none of the genes showed a significant rise in mRNA relative expression. When compared to the positive control group, the mRNA relative abundance of *EREG*, *PTGS2*, and *CTGF* genes at 12 h and *AREG* at 24 h was lower in groups treated with 0.3 μ M of both PAs, senecionine and seneciophylline ($P < 0.05$). Finally, 0.03 μ M senecionine reduced transcription levels of the *PTGS2* gene at 12 h, whereas 0.03 μ M seneciophylline reduced transcription levels of the same gene only at 24 h in comparison with positive control group ($P < 0.05$; **Figure 1**).




Figure 1 – Effect of Senecionine (A) or Seneciphylline (B) on expression of periovulatory EGF-Like cascade genes in bovine granulosa cells.

GC from large follicles (≥ 10 mm diameter) were cultured with serum for 24 h, and then in serum-free medium for another 18 h before addition of LH (100 ng/mL) and PAs (0, 0.03, and 0.3 μ M) for the times given. The Neg_Ctrl group was cultured without PAs and LH. Messenger RNA abundance was measured by real-time qPCR and normalized to the reference gene GAPDH. Bars represent the group mean $\pm$ SEM for three independent replicate cultures. Main effects of group, time, and their interaction were determined by ANOVA and are shown in the respective figure. Different letters indicate statistical significance among groups in the same time point ($P < 0.05$; Tukey-Kamer HSD).

3.4.2 PAs have no effect on cumulus cell expansion in in vitro matured COCs.

The expansion of cumulus cells was not blocked by any of the pyrrolizidine alkaloid concentrations evaluated. Indeed, there was no significant difference in the relative expansion of any group up to 6 hours of IVM, but from 12 hours of IVM to the end of the maturation period all groups showed a greater relative expansion than the negative control group ($P < 0.05$), although they did not differ from one another (*Figure 2*).




Figure 2 – Effect of graded concentrations (0, 0.03, and 0.3 μ M) of A) Senecionine or B) Seneciphylline on cumulus cells expansion.

COCs ($n=8$ /group/replicate) were individually matured and photographed at 0, 6, 12, and 24 h of IVM. The Neg_Ctrl group was cultured without PAs, FSH, and LH whereas Pos_Ctrl group was cultured without PAs. Relative cumulus expansion was assessed by dividing total surface area of each COC at a given time by the respective area at time 0. Points represent the group mean $\pm$ SEM for three independent replicate cultures. Main effects of group, time, and their interaction were determined by ANOVA and are shown in the respective figure. Different letters show statistically significant differences between groups in the same time point ($P < 0.05$; Tukey-Kamer HSD)

3.4.3 Effects of PAs added to the IVM medium on bovine embryo production and quality.

In this experiment 759 oocytes were in vitro matured with senecionine or seneciphylline at three different concentrations (0, 0.03, and 0.3 μ M). After that, they were fertilized and cultivated without pyrrolizidine alkaloids. Cleavage rates were assessed 48 hpi, and blastocyst rates were assessed at 168 hpi (*Table 2*).

Table 2 – Cleavage rates at 48 hpi and blastocyst rates at 168 hpi of COCs treated with graded concentrations of pyrrolizidine alkaloids during in vitro maturation.

Maturation treatment	IVM oocytes (N)	Cleavage		Blastocysts	
		(N)	(%)	(N)	(%)
Control	126	106	84.13	39	30.95
Senecionine 0.03 μ M	122	107	87.70	51	41.80
Senecionine 0.3 μ M	123	110	89.43	36	29.27
Control	129	103	79.84	41	31.78

Maturation treatment	IVM oocytes (N)	Cleavage		Blastocysts	
		(N)	(%)	(N)	(%)
Seneciophylline 0.03 μ M	132	109	82.58	57	43.18
Seneciophylline 0.3 μ M	127	109	85.83	47	37.01

Data represents three independent replicates.

Senecionine (**Figure 3A**) and seneciophylline (**Figure 3B**) had no effect on cleavage rates ($P>0.05$) in oocyte groups treated during maturation, when compared to control groups. Likewise, all maturation-treated groups had no differences in blastocyst developing rates ($P>0.05$) on day 7 after in vitro fertilization, when compared to the control groups (**Figure 3C; 3D**).




Figure 3 – Risk ratio for cleavage on day 2 (A; B) or blastocyst on day 7 (C; D) after in vitro fertilization of COCs treated with graded concentrations of pyrrolizidine alkaloids during in vitro maturation.

(A; C) Senecionine treated groups (0 μ M, 0.03 μ M, and 0.3 μ M). (B; D) Seneciophylline treated groups (0 μ M, 0.03 μ M, and 0.3 μ M). Dots represents RR and error bar represents 95% confidence interval of three independents replicates. RR: risk ratio. P: p-value.

Analyses of the transcription levels of genes linked to DNA repair, apoptosis, and energy metabolism were used to indirectly assess embryo quality.

Figure 4 shows the mRNA relative expression of some DNA double-strand repair related genes (*TP53BP1*, *RAD51*, and *XRCC6*) in blastocysts from groups treated with senecionine (**Figure 4A**) and seneciophylline (**Figure 4B**) during in vitro maturation. When compared to the control group, both 0.03 μ M and 0.3 μ M seneciophylline concentrations in the

maturation medium downregulated *TP53BP1* mRNA levels ($P < 0.05$). The addition of seneciphylline in the maturation medium had no effect on the quantity of transcripts from other genes involved in double-stranded DNA repair. In comparison to the control group, adding senecionine to the maturation medium has no effect on the transcription levels of any of the DNA double-strand repair related genes tested.




Figure 4 – Relative mRNA expression of genes associated with DNA double-strand repair in bovine embryos produced in vitro and treated with graded concentrations of pyrrolizidine alkaloids during in vitro maturation. A) Senecionine-treated groups (0, 0.03, and 0.3μM). B) Seneciphylline-treated groups (0, 0.03, and 0.3μM). The Control group was cultured without PAs. Messenger RNA abundance was measured by real-time qPCR and normalized to the reference gene GAPDH. Bars represent the group mean $\pm$ SEM for three independent replicate cultures. The effect of group was determined by ANOVA and is shown in the respective figure. Different letters indicate significant differences among groups ($P < 0.05$; Tukey-Kamer HSD).

There was no difference in mRNA expression for BAX, BCL2, or the BAX/BCL2 ratio between groups treated with senecionine or seneciphylline during in vitro maturation and groups treated with no pyrrolizidine alkaloids (**Figure 5**).




Figure 5 – Relative mRNA expression of BAX, BCL2 and BAX/BCL2 ratio in bovine embryos produced in vitro and treated with graded concentrations of pyrrolizidine alkaloids during in vitro maturation.

A) Senecionine-treated groups (0, 0.03, and 0.3 μ M). B) Seneciophylline-treated groups (0, 0.03, and 0.3 μ M). The Control group was cultured without PAs. Messenger RNA abundance was measured by real-time qPCR and normalized to the reference gene GAPDH. Bars represent the group mean $\pm$ SEM for three independent replicate cultures. The effect of group was determined by ANOVA and is shown in the respective figure. Different letters indicate significant differences among groups ($P < 0.05$; Tukey-Kamer HSD).

The relative mRNA abundance of GLUT3 gene was lower in blastocysts from the group treated with 0.3 μ M of senecionine (**Figure 6A**) in the maturation medium ($P < 0.05$) but neither 0.03 μ M of senecionine, nor seneciophylline (**Figure 6B**) modulated mRNA expression of the GLUT1 or GLUT3 genes, when compared to their respective control groups.




Figure 6 – Relative mRNA expression of GLUT1 and GLUT3 in bovine embryos produced in vitro and treated with graded concentrations of pyrrolizidine alkaloids during in vitro maturation.

A) Senecionine-treated groups (0, 0.03, and 0.3 μ M). B) Seneciophylline-treated groups (0, 0.03, and 0.3 μ M). The Control group was cultured without PAs. Messenger RNA abundance was measured by real-time qPCR and normalized to the reference gene GAPDH. Bars represent the group mean $\pm$ SEM for three independent replicate cultures. The effect of group was determined by ANOVA and is shown in the respective figure. Different letters indicate significant differences among groups ($P < 0.05$; Tukey-Kamer HSD).

3.4.4 Effects of PAs added to the IVC medium on bovine embryo production and quality.

In this experiment, 1244 oocytes were in vitro matured and fertilized without pyrrolizidine alkaloids. They were then in vitro cultured at three different senecionine or seneciophylline concentrations (0, 0.03, and 0.3 μ M) beginning at 18–20 hpi. Cleavage and blastocyst rates were assessed at 48 and 168 hpi, respectively. (**Table 3**).

Table 3 – Cleavage rates at 48 hpi and blastocyst rates at 168 hpi of COCs treated with graded concentrations of pyrrolizidine alkaloids added to IVC medium.

Maturation treatment	IVM oocytes (N)	Cleavage		Blastocysts	
		(N)	(%)	(N)	(%)
Control	209	164	78.47	63	30.14
Senecionine 0.03 μ M	203	135	65.50	43	21.18
Senecionine 0.3 μ M	195	118	60.51	27	13.85
Control	215	168	78.14	66	30.70
Seneciophylline 0.03 μ M	212	137	64.62	40	18.87
Seneciophylline 0.3 μ M	210	129	61.43	34	16.19

Data represents five replicates.

When pyrrolizidine alkaloids were added to the culture media at 18-20 hpi all of the treated groups had a lower rate of cleavage than the control groups ($P < 0.05$). The cleavage

risk ratio with 0,03 μM senecionine was 0.848 [0.751 – 0.956; 95%IC] (**Figure 7A**). With 0.03 μM seneciphylline, the cleavage risk ratio was 0.827 [0.731 – 0.934; 95% IC] (**Figure 7B**). This effect was more pronounced when the highest concentrations of senecionine or seneciphylline were utilized. The cleavage risk ratio with 0,3 μM senecionine was 0.771 [0.674 – 0.881; 95%IC] (**Figure 7A**). With 0.3 μM seneciphylline, the cleavage risk ratio was 0.786 [0.691 – 0.893; 95% IC] (**Figure 7B**).

When senecionine or seneciphylline were added to the culture media 18-20 hpi, blastocyst rates on day 7 after fertilization were lowered in an even more pronounced way than the cleavage rate, as compared to groups cultured without pyrrolizidine alkaloids ($P < 0.05$). The blastocyst risk ratio with 0,03 μM senecionine was 0.703 [0.502 – 0.983; 95%IC] (**Figure 7C**). With 0.03 μM seneciphylline, the blastocyst risk ratio was 0.615 [0.435 – 0.866; 95% IC] (**Figure 7D**). Here again, this effect was more pronounced when the highest concentrations of senecionine or seneciphylline were utilized. The cleavage risk ratio with 0,3 μM senecionine was 0.459 [0.305 – 0.689; 95%IC] (**Figure 7C**). With 0.3 μM seneciphylline, the cleavage risk ratio was 0.527 [0.365 – 0.761; 95% IC] (**Figure 7D**).




Figure 7 – Risk ratio for cleavage on day 2 (A; B) or blastocyst on day 7 (C; D) after in vitro fertilization of zygotes treated with graded concentrations of pyrrolizidine alkaloids added to the medium at 18-20 hpi.

(A; C) Senecionine treated groups (0, 0.03, and 0.3 μM). (B; D) Seneciphylline treated groups (0, 0.03, and 0.3 μM). Dots represents RR and error bar represents 95% confidence interval of five independent replicates. RR: risk ratio. P: p-value.

In this experiment, embryo quality was assessed using the same genes that were evaluated in the prior experiment.

Figure 8 shows the mRNA relative expression of some DNA double-strand repair related genes (*TP53BP1*, *RAD51*, and *XRCC6*) in blastocysts from groups treated with senecionine (**Figure 8A**) and seneciphylline (**Figure 8B**) 18-20 hpi. When compared to the control group, 0.3 μ M seneciphylline concentration in the IVC medium, but not 0.03 μ M seneciphylline concentration, upregulates *RAD51* mRNA levels ($P < 0.05$). The addition of seneciphylline in the IVC medium had no effect on the quantity of transcripts from other genes involved in double-stranded DNA repair. In comparison to the control group, adding senecionine to the IVC medium has no effect on the transcription levels of any of the DNA double-strand repair related genes tested.




Figure 8 – Relative mRNA expression of genes associated with DNA double-strand repair in bovine in vitro produced blastocysts treated with graded concentrations of pyrrolizidine alkaloids added to the medium at 18-20 hpi.

A) Senecionine-treated groups (0, 0.03, and 0.3 μ M). B) Seneciphylline-treated groups (0, 0.03, and 0.3 μ M). The Control group was cultured without PAs. Messenger RNA abundance was measured by real-time qPCR and normalized to the reference gene GAPDH. Bars represent the group mean $\pm$ SEM for five independent replicate cultures. The effect of group was determined by ANOVA and is shown in the respective figure. Different letters indicate significant differences among groups ($P < 0.05$; Tukey-Kramer HSD).

Figure 9 shows the mRNA relative expression of *BAX* and *BCL2* genes and the *BAX/BCL2* ratio in blastocysts from groups treated with senecionine (**Figure 9A**) and seneciphylline (**Figure 9B**) 18-20 hpi. When compared to the control group, 0.3 μM senecionine concentration in the IVC medium downregulates the anti-apoptotic gene *BCL2* ($P < 0.05$), but 0.03 μM senecionine had no effect on the same gene expression. Senecionine showed no effect on *BAX* gene transcript levels or on the *BAX/BCL2* ratio when added to the IVC medium. Adding seneciphylline to the IVC medium had no influence on the transcription levels of *BAX* or *BCL2* genes, nor on the *BAX/BCL2* ratio, as compared to their respective control groups.




Figure 9 – Relative mRNA expression of *BAX*, *BCL2*, and *bax/bcl2* ratio in bovine in vitro produced blastocysts treated with graded concentrations of pyrrolizidine alkaloids added to the medium at 18-20 hpi. A) Senecionine-treated groups (0, 0.03, and 0.3 μM). B) Seneciphylline-treated groups (0, 0.03, and 0.3 μM). The Control group was cultured without PAs. Messenger RNA abundance was measured by real-time qPCR and normalized to the reference gene GAPDH. Bars represent the group mean $\pm$ SEM for five independent replicate cultures. The effect of group was determined by ANOVA and is shown in the respective figure. Different letters indicate significant differences among groups ($P < 0.05$; Tukey-Kamer HSD).

There was no difference in mRNA expression of the *GLUT1* or *GLUT3* genes between groups treated with senecionine or seneciphylline after denudation and groups treated with no pyrrolizidine alkaloids, as shown in **Figure 10**.




Figure 10– Relative mRNA expression of GLUT1 and GLUT3 in bovine in vitro produced blastocysts treated with graded concentrations of pyrrolizidine alkaloids added to the medium at 18-20 hpi.

A) Senecionine-treated groups (0, 0.03, and 0.3 μ M). B) Seneciphylline-treated groups (0, 0.03, and 0.3 μ M). The Control group was cultured without PAs. Messenger RNA abundance was measured by real-time qPCR and normalized to the reference gene GAPDH. Bars represent the group mean $\pm$ SEM for five independent replicate cultures. The effect of group was determined by ANOVA and is shown in the respective figure. Different letters indicate significant differences among groups ($P < 0.05$; Tukey-Kamer HSD).

3.5 DISCUSSION

This study tested the hypothesis that low concentrations of pyrrolizidine alkaloids, insufficient to cause chronic seneciosis, could act directly on ovarian cells and/or embryos, affecting the periovulatory Epidermal Growth Factor-Like (EGF-Like) cascade and early embryonic development.

Our significant findings are: (1) Senecionine as well as seneciphylline downregulates expression of periovulatory EGF-Like cascade genes in bovine granulosa cells in a time and dose dependent manner; (2) cumulus cells expansion was not blocked by any of the pyrrolizidine alkaloid concentrations evaluated; (3) when added to maturation medium PAs don't alter embryo production, however both cleavage and blastocyst rates decrease when senecionine or seneciphylline were added to IVC medium; (4) PAs downregulates the transcript level of genes related to embryonic energy metabolism, DNA repair and apoptosis in blastocysts, suggesting that blastocysts that had contact with PAs have lower developmental competence; and (5) low concentrations of both senecionine and

seneciophylline can act directly on bovine granulosa cells and embryos, disrupting some reproductive processes in vitro.

Ovulation is a complex process that is initiated by the luteinizing hormone surge and is controlled by the temporal and spatial expression of specific genes. LH triggers a cascade of signaling processes that spread throughout the preovulatory follicle, culminating in the ovarian follicle wall rupture and the release of an oocyte into the oviduct for fertilization [28,29]. This cascade, however, requires the activation of EGF receptor (EGFR). When the EGFR is activated, the cumulus expresses *AREG* and *EREG* and increase *PTGS2* levels. In mural granulosa cells, the similar loop occurs, with *AREG* and *EREG* working in an autocrine manner to intensify the cascade [21,30–32]. It has been shown that the transcriptional activity of Yes-associated protein (*YAP*), an effector of the Hippo pathway, regulates the EGF signaling cascade in bovine mural granulosa and cumulus cells and directly interferes with the processes of cumulus expansion and ovulation that are triggered by LH [22,33]. In this study, we used a previously described granulosa cell culture system [21,22] in which abundance of mRNA encoding the EGF-like factors *EREG* and *AREG*, and other critical preovulatory genes is acutely upregulated by LH. Our results clearly indicate that inclusion of 0.3 μ M of senecionine or 0.3 μ M of seneciophylline to the culture medium effectively blocked the effect of LH on *EREG* and *CTGF*, mRNA abundance at 12 hours, on *PTGS2* transcripts from 12 hours, and *AREG* mRNA abundance at 24 hours. Similarly, *PTGS2* mRNA abundance was blocked with 0.03 μ M of both PAs. In mice, disruption of the EGF-Like cascade impairs ovulation, indicating that the activation of this pathway is essential for LH-induced ovulation to occur [34]. The importance of *PTGS2* in ovulation also has been demonstrated. *PTGS2*-null mice failed to ovulate [35] whereas in cattle, the intrafollicular injection of a *PTGS2*-selective inhibitor inhibited ovulation [36]. In the same way, it has been demonstrated that *YAP* signaling is critical for expression of EGFR and downstream target genes in bovine granulosa cells in vitro [22]. Overall, our findings suggest that even low concentrations of senecionine and seneciophylline can impair ovulation in cattle.

In most mammals, before ovulation, cumulus cells synthesize a large amount of the polysaccharide hyaluronan (HA), which is organized into a highly hydrated muco-elastic matrix. This process is called cumulus expansion or mucification [37]. In vitro, the remotion of COC from the follicle is sufficient to lead to cumulus expansion, whereas in vivo, this process is triggered by LH surge. Cumulus expansion is dependent on the expression of EGF-like ligands, *PTGS2*, and *YAP* in granulosa cells, however these genes are also expressed in

cumulus cells when stimulated by FSH [33,38]. Intriguingly, none of the pyrrolizidine alkaloid concentrations tested in our experiment prevented cumulus cells expansion. These findings, indeed, contradict what we found in the granulosa cell experiment, in which we demonstrated that PAs inhibit the key genes for cumulus expansion. We suppose that this discrepancy could be due to the in vitro cumulus expansion models used herein, which employ supraphysiological levels of FSH, LH, and fetal calf serum during the entire course of the experiment, and that may be masking the effects of PAs.

The ability of a blastocyst to establish a pregnancy and generate a living offspring is the strongest indicator of its quality. Because it is impossible to transfer every embryo, the achievement of the blastocyst stage and morphological appearance are commonly employed as measures of oocyte developmental capacity. For the same reason, other ways of determining the quality of a blastocyst are used, such as cell counts, cryotolerance, and gene expression patterns [39,40]. In this study, gene expression analyses were used to assess embryo quality.

Cell repair machinery is responsible for protecting the genome from endogenous and exogenous effects that induce DNA damage, furthermore DNA repair during the early stages of embryonic development has one of the most significant effects on embryonic fate [41]. DNA double-strand breaks (DSB) are the most deleterious type of damage and can substantially alter genetic integrity [42]. The presence of DSBs can alter the kinetics of embryo cell cleavage and development to the blastocyst stage, therefore only embryos with less DNA damage and/or superior capacity for DNA repair are able to achieve the blastocyst stage [43]. DNA repair and cell cycle checkpoint genes are up-regulated in late-cleaving embryos [43,44].

BCL2 family members regulate the intrinsic apoptotic pathway and may be considered as good markers of developmental potential. Thus, the balance between pro-apoptotic and anti-apoptotic factors can determine whether and when a cell becomes apoptotic. BCL2 protein is considered an anti-apoptotic agent whereas BAX protein acts as pro apoptotic [45]. The expression of *BCL2* gene is increased in bovine and pig embryos with good morphological quality [46,47] and in non-fragmented mouse blastocysts [45].

Glucose transporters (GLUT) carry glucose through COCs and embryos in a passive, energy-independent manner during culture [46]. *GLUT-3* is one of the key isoforms of glucose transporters and, in conjunction with *GLUT-1* are the main isoform during bovine early embryo development. The association between a faster growth rate and higher rates of

expression of *GLUT-1* and *GLUT-3* genes, may enhance embryo metabolism and developmental competence [48]. Embryos with a higher relative abundance of *GLUT-3* compared to their counterparts has significantly high levels of development to the blastocyst stage [49].

In our experiments only four of evaluated genes were affected by senecionine or seneciphylline. When PAs were added to the maturation medium, there were no differences in cleavage or blastocyst rate, but the senecionine-treated group (0.3 μ M) showed lower blastocyst expression of *GLUT-3* mRNA, implying that these blastocysts are less competent. On oocytes matured with Seneciphylline (0.3 and 0.03 μ M), blastocyst *TP53BP1* mRNA abundance decreases during maturation. The cellular response to DSB is regulated by *TP53BP1*, which directs repair to the NHEJ pathway. The *BRCA1* modulator, on the other hand, increases *TP53BP1* dephosphorylation throughout the S/G2 phases of the cell cycle, directing repair to the HR pathway. Seneciphylline (0.3 μ M) in IVC medium up-regulated the expression of *RAD51*, another integrant of the HR pathway [50]. One of the characteristics of PAs is their ability to form adducts with proteins and nucleic acids [51,52]. These findings suggest that seneciphylline may have caused DNA damage to the embryo. Finally, adding 0.3 μ M senecionine to the IVC medium reduced the amount of *BCL2* transcripts. This means that there may be an imbalance between the regulators of the intrinsic apoptosis pathway and that anti-apoptotic mechanisms may not operate effectively, favoring apoptosis.

In conventional in vitro bovine embryo production laboratory, the cleavage rate ranges between 75% and 90%, whereas the blastocyst rate ranges between 30% and 40% [25,39,44]. In this study, the cleavage rate of the control groups ranged from 78.14% to 84.13%, whereas the blastocyst rate ranged from 30.14% to 31.78%. Both PAs, senecionine and seneciphylline, decreased cleavage and blastocyst rate in treated groups when added to IVC medium, but not during maturation. As a result, the presumptive zygote cultured in 0.03 μ M senecionine had 15.2% less risk of cleavage and 29.7% less risk of developing into a blastocyst when compared to control group. In its turn, the putative zygote cultured in 0.3 μ M senecionine had 22.9% less risk of cleavage and 54.1% less risk of developing into a blastocyst when compared to control group. When the presumptive zygote was cultured with 0.03 μ M seneciphylline, it had 17.3% less risk of cleavage and 38.5% less risk of becoming a blastocyst, compared to the control group. Finally, the presumptive zygote cultured in 0.3 μ M seneciphylline had 21.4% less risk of cleavage and 47.3% less risk of becoming a blastocyst, compared to the control group. The results of the genic expression and the embryonic

production, taken together, suggest that the blastocysts formed in the treated groups are not only less numerous, but also have a lesser developmental capacity than their control counterparts.

Despite variability in pyrrolizidine alkaloids sensitivity between species or even between tissues, cytochrome P450 bioactivation is needed for toxicity. Therefore, in models where cells are unable to bioactivate PAs, exogenous pre metabolism is required to observe the effects of the PAs. Consequently, without metabolism, no relevant effects were observed up to the highest tested concentrations (500 μ M) in human HUVEC, whereas cells incubated with PA after preincubation using liver homogenate showed treatment-induced responses, both cytotoxicity (300 μ M) and alterations of prostanoid synthesis (200 μ M) [53]. In the same way, in vitro experiments showed no cytotoxicity to freshly isolated mouse LSECs up to 500 μ M senecionine. However, metabolic activation of senecionine by preincubation with primary mouse hepatocytes increased the cytotoxicity to cultivated LSECs with an EC_{50} of approximately 22 μ M [54]. Senecionine and seneciphylline were found to have a detrimental effect on the in vitro embryo quality and production and on granulosa cell culture groups in this investigation. These findings show that APs are metabolically activated by granulosa and embryo cells, which suggest that they can directly affects these cells even if no hepatic metabolism takes place.

In summary, this study demonstrates that both senecionine and seneciphylline have significant toxicological effects in a bovine granulosa cell model as well as in vitro produced bovine embryos, suggesting that pyrrolizidine alkaloids, even at low concentrations, have the potential to directly affect cattle reproductive physiology.

Source of Funding

Conflict of interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

The authors would like to thank El'Golli and Verdi abattoirs for providing the bovine ovaries.

3.6 REFERENCES

- [1] C.S.L. Barros, L.M.L. Castilhos, D.R. Rissi, G.D. Kommers, R.R. Rech, Liver biopsy for the diagnosis of *Senecio brasiliensis* (Asteraceae) poisoning in cattle, Pesqui.

- 583 Veterinária Bras. 27 (2007) 53–60. <https://doi.org/10.1590/S0100->
584 736X2007000100010.
- 585 [2] F. Riet-Correa, R.M.T. Medeiros, Intoxicações por plantas em ruminantes no Brasil e
586 no Uruguai: importância econômica, controle e riscos para a saúde pública, Pesqui.
587 Veterinária Bras. 21 (2001) 38–42. <https://doi.org/10.1590/S0100->
588 736X2001000100008.
- 589 [3] C.H. Tokarnia, J. Döbereiner, P.V. Peixoto, Poisonous plants affecting livestock in
590 Brazil, Toxicon. 40 (2002) 1635–1660.
- 591 [4] T.M. Sandini, M. Sayuri, U. Berto, Senecio brasiliensis e alcaloides pirrolizidínicos:
592 toxicidade em animais e na saúde humana, Rev. Biotemas Biotemas. 26 (2013) 83–92.
593 <https://doi.org/10.5007/2175-7925.2013v26n2p83>.
- 594 [5] D.R. Rissi, R.R. Rech, F. Pierezan, A.L. Gabriel, M.E. Trost, J.S. Brum, G.D.
595 Kommers, C.S.L. Barros, Plant and plant-associated mycotoxins poisoning in cattle in
596 Rio Grande do Sul, Brazil: 461 cases, Pesqui. Veterinária Bras. 27 (2007) 261–268.
- 597 [6] F.C. Karam, A.L. Schild, J.R.B. Mello, Poisoning by Senecio spp. in cattle in southern
598 Brazil: favorable conditions and control measures, Pesqui. Veterinária Bras. 31 (2011)
599 603–609. <https://doi.org/10.1590/S0100-736X2011000700010>.
- 600 [7] T. Hartmann, L. Witte, Chemistry, Biology and Chemoecology of the Pyrrolizidine
601 Alkaloids, in: S.W. Pelletier (Ed.), Alkaloids Chem. Biol. Perspect., Pergamon,
602 Oxford, 1995: pp. 155–233. <https://doi.org/10.1016/B978-0-08-042089-9.50011-5>.
- 603 [8] J.R. Trigo, I.R. Leal, N.I. Matzenbacher, T.M. Lewinsohn, Chemotaxonomic value of
604 pyrrolizidine alkaloids in southern Brazil Senecio (Senecioneae: Asteraceae), Biochem.
605 Syst. Ecol. 31 (2003) 1011–1022. [https://doi.org/10.1016/S0305-1978\(03\)00038-3](https://doi.org/10.1016/S0305-1978(03)00038-3).
- 606 [9] R. Adams, M. Gianturco, Senecio alkaloids: the alkaloids of Senecio brasiliensis,
607 fremonti and ambrosioides, J. Am. Chem. Soc. 78 (1956) 5315–5317.
608 <https://doi.org/10.1021/ja01601a044>.
- 609 [10] A.S. Prakash, T.N. Pereira, P.E.. Reilly, A.A. Seawright, Pyrrolizidine alkaloids in
610 human diet, Mutat. Res. Toxicol. Environ. Mutagen. 443 (1999) 53–67.
611 [https://doi.org/10.1016/S1383-5742\(99\)00010-1](https://doi.org/10.1016/S1383-5742(99)00010-1).
- 612 [11] P.R. Cheeke, Toxicity and metabolism of pyrrolizidine alkaloids, J. Anim. Sci. 66
613 (1988) 2343–2350. <https://doi.org/10.2527/jas1988.6692343x>.
- 614 [12] D.N. Skilleter, A.R. Mattocks, G.E. Neal, Sensitivity of different phases of the cell
615 cycle to selected hepatotoxins in cultured liver-derived (BL9L) cells, Xenobiotica. 18
616 (1988) 699–705. <https://doi.org/10.3109/00498258809041708>.
- 617 [13] J.R. Basile, J.M.F. Diniz, W. Okano, S.M. Cirio, L.C. Leite, Intoxicação por Senecio
618 spp. (Compositae) em bovinos no sul do Brasil, Acta Sci. Vet. 33 (2005) 63–68.
- 619 [14] C.H. Tokarnia, J. Dobereiner, Intoxicação experimental por Senecio brasiliensis
620 (Compositae) em bovinos, Pesqui. Vet. Bras. 4 (1984) 39–65.
- 621 [15] U. Candrian, U. Zweifel, J. Luethy, C. Schlatter, Transfer of orally administered 3H-
622 seneciphylline into cow's milk, J. Agric. Food Chem. 39 (1991) 930–933.
623 <https://doi.org/10.1021/jf00005a026>.
- 624 [16] F.S.C. Karam, M.P. Soares, M. Haraguchi, F. Riet-correa, M. del C. Méndez, J.A.
625 Jarenkow, Epidemiological aspects of seneciosis in southern Rio Grande do Sul, Brazil,

- 626 Pesqui. Veterinária Bras. 24 (2004) 191–198.
- 627 [17] R.B. Lucena, D.R. Rissi, L.A. Maia, M.M. Flores, A.F.M. Dantas, V.M. da T. Nobre,
628 F. Riet-Correa, C.S.L. Barros, Poisoning by pyrrolizidine alkaloids in ruminants and
629 horses in Brazil, Pesqui. Veterinária Bras. 30 (2010) 447–452.
630 <https://doi.org/10.1590/S0100-736X2010000500013>.
- 631 [18] J.K. Leighton, S. Canning, H.D. Guthrie, J.M. Hammond, Expression of cytochrome
632 P450 1A1, an estrogen hydroxylase, in ovarian granulosa cells is developmentally
633 regulated, J. Steroid Biochem. Mol. Biol. 52 (1995) 351–356.
634 [https://doi.org/10.1016/0960-0760\(94\)00185-O](https://doi.org/10.1016/0960-0760(94)00185-O).
- 635 [19] A.A. Krabbe, M.A. Gonçalves, R. Pozzobon, C.R.M. Pessoa, M.P. Soares, R.A. Costa,
636 B.L. Anjos, A.A. Krabbe, M.A. Gonçalves, R. Pozzobon, C.R.M. Pessoa, M.P. Soares,
637 R.A. Costa, B.L. Anjos, Megalocitose de células luteínicas grandes de vacas prenhes
638 com seneciose crônica, Pesqui. Veterinária Bras. 35 (2015) 33–38.
639 <https://doi.org/10.1590/S0100-736X2015000100008>.
- 640 [20] T.M. Sandini, M.S.B. Udo, T.M. Reis-Silva, M.M. Bernardi, H. de S. Spinosa, Prenatal
641 exposure to integerrimine N-oxide impaired the maternal care and the physical and
642 behavioral development of offspring rats, Int. J. Dev. Neurosci. 36 (2014) 53–63.
643 <https://doi.org/10.1016/j.ijdevneu.2014.05.007>.
- 644 [21] V.M. Portela, G. Zamberlam, P.B.D. Gonçalves, J.F.C. de Oliveira, C.A. Price, Role of
645 Angiotensin II in the Perioovulatory Epidermal Growth Factor-Like Cascade in Bovine
646 Granulosa Cells In Vitro¹, Biol. Reprod. 85 (2011) 1167–1174.
647 <https://doi.org/10.1095/biolreprod.111.094193>.
- 648 [22] E.C. Dos Santos, A. Lalonde-Larue, A.Q. Antoniazzi, M.H. Barreta, C.A. Price, P.B.
649 Dias Gonçalves, V.M. Portela, G. Zamberlam, YAP signaling in preovulatory
650 granulosa cells is critical for the functioning of the EGF network during ovulation,
651 Mol. Cell. Endocrinol. 541 (2022) 111524. <https://doi.org/10.1016/j.mce.2021.111524>.
- 652 [23] L. Leibfried, N.L. First, Characterization of bovine follicular oocytes and their ability
653 to mature, J. Anim. Sci. 48 (1979) 76–86. <https://doi.org/10.2527/jas1979.48176x>.
- 654 [24] J.J. Parrish, J. Susko-Parrish, M.A. Winer, N.L. First, Capacitation of bovine sperm by
655 heparin, Biol. Reprod. 38 (1988) 1171–1180.
656 <https://doi.org/10.1095/biolreprod38.5.1171>.
- 657 [25] P. Holm, P.J. Booth, M.H. Schmidt, T. Greve, H. Callesen, High bovine blastocyst
658 development in a static in vitro production system using sofaa medium supplemented
659 with sodium citrate and myo-inositol with or without serum-proteins, Theriogenology.
660 52 (1999) 683–700. [https://doi.org/10.1016/S0093-691X\(99\)00162-4](https://doi.org/10.1016/S0093-691X(99)00162-4).
- 661 [26] J. Vandesompele, K. De Preter, F. Pattyn, B. Poppe, N. Van Roy, A. De Paepe, F.
662 Speleman, Accurate normalization of real-time quantitative RT-PCR data by geometric
663 averaging of multiple internal control genes, Genome Biol. 3 (2002) 1–12.
664 <https://doi.org/https://doi.org/10.1186/gb-2002-3-7-research0034>.
- 665 [27] M.W. Pfaffl, A new mathematical model for relative quantification in real-time RT-
666 PCR, Nucleic Acids Res. 29 (2001) 2002–2007. <https://doi.org/10.1093/nar/29.9.e45>.
- 667 [28] J.S. Richards, D.L. Russell, S. Ochsner, L.L. Espey, Ovulation : new dimensions and
668 new regulators of the inflammatory-like response, Annu. Rev. Physiol. 64 (2002) 69–
669 92. <https://doi.org/10.1146/annurev.physiol.64.081501.131029>.

- [29] W.W. Thatcher, A 100-year review: historical development of female reproductive physiology in dairy cattle, *J. Dairy Sci.* 100 (2017) 10272–10291. <https://doi.org/10.3168/jds.2017-13399>.
- [30] J.-Y. Park, Y.-Q. Su, M. Ariga, E. Law, S.-L.C. Jin, M. Conti, EGF-Like growth factors as mediators of LH action in the ovulatory follicle, *Science* (80-.). 303 (2004) 682–684. <https://doi.org/10.1126/science.1092463>.
- [31] M. Shimada, I. Hernandez-Gonzalez, I. Gonzalez-Robayna, J.S. Richards, Paracrine and Autocrine Regulation of Epidermal Growth Factor-Like Factors in Cumulus Oocyte Complexes and Granulosa Cells: Key Roles for Prostaglandin Synthase 2 and Progesterone Receptor, *Mol. Endocrinol.* 20 (2006) 1352–1365. <https://doi.org/10.1210/me.2005-0504>.
- [32] S. Panigone, M. Hsieh, M. Fu, L. Persani, M. Conti, Luteinizing Hormone Signaling in Preovulatory Follicles Involves Early Activation of the Epidermal Growth Factor Receptor Pathway, *Mol. Endocrinol.* 22 (2008) 924–936. <https://doi.org/10.1210/me.2007-0246>.
- [33] J. Koch, V.M. Portela, E.C. dos Santos, D. Missio, L.G. de Andrade, Z. da Silva, B.G. Gasperin, A.Q. Antoniazzi, P.B.D. Gonçalves, G. Zamberlam, The Hippo pathway effectors YAP and TAZ interact with EGF-like signaling to regulate expansion-related events in bovine cumulus cells in vitro, *J. Assist. Reprod. Genet.* (2022). <https://doi.org/10.1007/s10815-021-02384-x>.
- [34] M. Hsieh, D. Lee, S. Panigone, K. Horner, R. Chen, A. Theologis, D.C. Lee, D.W. Threadgill, M. Conti, Luteinizing Hormone-dependent activation of the epidermal growth factor network is essential for ovulation, *Mol. Cell. Biol.* 27 (2007) 1914–1924. <https://doi.org/10.1128/MCB.01919-06>.
- [35] B.J. Davis, D.E. Lennard, C.A. Lee, H.F. Tiano, S.G. Morham, W.C. Wetsel, R. Langenbach, Anovulation in Cyclooxygenase-2-Deficient Mice Is Restored by Prostaglandin E2 and Interleukin-1 β , *Endocrinology*. 140 (1999) 2685–2695. <https://doi.org/10.1210/endo.140.6.6715>.
- [36] M.W. Peters, J.R. Pursley, G.W. Smith, Inhibition of intrafollicular PGE2 synthesis and ovulation following ultrasound-mediated intrafollicular injection of the selective cyclooxygenase-2 inhibitor NS-398 in cattle, *J. Anim. Sci.* 82 (2004) 1656–1662. <https://doi.org/10.2527/2004.8261656x>.
- [37] E. Nagyova, A. Camaioni, R. Prochazka, A.J. Day, A. Salustri, Synthesis of Tumor Necrosis Factor Alpha-Induced Protein 6 in Porcine Preovulatory Follicles: A Study with A38 Antibody, *Biol. Reprod.* 78 (2008) 903–909. <https://doi.org/10.1095/biolreprod.107.064832>.
- [38] E.S. Caixeta, M.F. Machado, P. Ripamonte, C. Price, J. Buratini, Effects of FSH on the expression of receptors for oocyte-secreted factors and members of the EGF-like family during in vitro maturation in cattle, *Reprod. Fertil. Dev.* 25 (2013) 890–899. <https://doi.org/10.1071/RD12125>.
- [39] P. Lonergan, D. Rizos, A. Gutierrez-Adan, T. Fair, M. Boland, Oocyte and Embryo Quality: Effect of Origin, Culture Conditions and Gene Expression Patterns, *Reprod. Domest. Anim.* 38 (2003) 259–267. <https://doi.org/10.1046/j.1439-0531.2003.00437.x>.
- [40] J.L. Leroy, G. Opsomer, S. De Vlieghe, T. Vanholder, L. Goossens, A. Geldhof, P.E. Bols, A. de Kruif, A. Van Soom, Comparison of embryo quality in high-yielding dairy

- cows, in dairy heifers and in beef cows, *Theriogenology*. 64 (2005) 2022–2036. <https://doi.org/10.1016/j.theriogenology.2005.05.003>.
- [41] E. V. Khokhlova, Z.S. Fesenko, J. V. Sopova, E.I. Leonova, Features of DNA Repair in the Early Stages of Mammalian Embryonic Development, *Genes (Basel)*. 11 (2020) 1138. <https://doi.org/10.3390/genes11101138>.
- [42] S. Titus, F. Li, R. Stobezki, K. Akula, E. Unsal, K. Jeong, M. Dickler, M. Robson, F. Moy, S. Goswami, K. Oktay, Impairment of BRCA1-Related DNA Double-Strand Break Repair Leads to Ovarian Aging in Mice and Humans, *Sci. Transl. Med.* 5 (2013). <https://doi.org/10.1126/scitranslmed.3004925>.
- [43] R.C. Bohrer, A.R.S. Coutinho, R. Duggavathi, V. Bordinon, The incidence of DNA double-strand breaks is higher in late-cleaving and less developmentally competent porcine embryos, *Biol. Reprod.* 93 (2015) 1–8. <https://doi.org/10.1095/biolreprod.115.130542>.
- [44] M.H. Barreta, B.G. Gasperin, V.B. Rissi, M.P. de Cesaro, R. Ferreira, J.F. de Oliveira, P.B.D. Gonçalves, V. Bordinon, Homologous recombination and non-homologous end-joining repair pathways in bovine embryos with different developmental competence, *Exp. Cell Res.* 318 (2012) 2049–2058. <https://doi.org/10.1016/j.yexcr.2012.06.003>.
- [45] D. Boruszewska, E. Sinderewicz, I. Kowalczyk-Zieba, K. Grycmacher, I. Woclawek-Potocka, The effect of lysophosphatidic acid during in vitro maturation of bovine cumulus–oocyte complexes: cumulus expansion, glucose metabolism and expression of genes involved in the ovulatory cascade, oocyte and blastocyst competence, *Reprod. Biol. Endocrinol.* 13 (2015) 44. <https://doi.org/10.1186/s12958-015-0044-x>.
- [46] P. Ramos-Ibeas, I. Gimeno, K. Cañón-Beltrán, A. Gutiérrez-Adán, D. Rizo, E. Gómez, Senescence and Apoptosis During in vitro Embryo Development in a Bovine Model, *Front. Cell Dev. Biol.* 8 (2020). <https://doi.org/10.3389/fcell.2020.619902>.
- [47] M.Y. Yang, R. Rajamahendran, Involvement of Apoptosis in the Atresia of Nonovulatory Dominant Follicle During the Bovine Estrous Cycle, *Biol. Reprod.* 63 (2000) 1313–1321. <https://doi.org/10.1095/biolreprod63.5.1313>.
- [48] M. Bertolini, S.W. Beam, H. Shim, L.R. Bertolini, A.L. Moyer, T.R. Famula, G.B. Anderson, Growth, development, and gene expression by in vivo- and in vitro-produced day 7 and 16 bovine embryos, *Mol. Reprod. Dev.* 63 (2002) 318–328. <https://doi.org/10.1002/mrd.90015>.
- [49] K.M. Morton, D. Herrmann, B. Sieg, C. Struckmann, W.M.C. Maxwell, D. Rath, G. Evans, A. Lucas-Hahn, H. Niemann, C. Wrenzycki, Altered mRNA expression patterns in bovine blastocysts after fertilisation in vitro using flow-cytometrically sex-sorted sperm, *Mol. Reprod. Dev.* 74 (2007) 931–940. <https://doi.org/10.1002/mrd.20573>.
- [50] F. Marini, C.C. Rawal, G. Liberi, A. Pellicioli, Regulation of DNA double strand breaks processing: focus on barriers, *Front. Mol. Biosci.* 6 (2019). <https://doi.org/10.3389/fmolb.2019.00055>.
- [51] R.J. Molyneux, D.L. Gardner, S.M. Colegate, J.A. Edgar, Pyrrolizidine alkaloid toxicity in livestock: a paradigm for human poisoning?, *Food Addit. Contam. Part A*. 28 (2011) 293–307. <https://doi.org/10.1080/19440049.2010.547519>.
- [52] P.P. Fu, Q. Xia, G. Lin, M.W. Chou, Pyrrolizidine alkaloids: genotoxicity, metabolism

- 759 enzymes, metabolic activation and mechanisms, *Drug Metab. Rev.* 36 (2004) 1–55.
760 <https://doi.org/10.1081/DMR-120028426>.
- 761 [53] J. Ebmeyer, J. Behrend, M. Lorenz, G. Günther, R. Reif, J.G. Hengstler, A. Braeuning,
762 A. Lampen, S. Hessel-Pras, Pyrrolizidine alkaloid-induced alterations of prostanoid
763 synthesis in human endothelial cells, *Chem. Biol. Interact.* 298 (2019) 104–111.
764 <https://doi.org/10.1016/j.cbi.2018.11.007>.
- 765 [54] S. Hessel-Pras, A. Braeuning, G. Guenther, A. Adawy, A.M. Enge, J. Ebmeyer, C.J.
766 Henderson, J.G. Hengstler, A. Lampen, R. Reif, The pyrrolizidine alkaloid senecionine
767 induces CYP-dependent destruction of sinusoidal endothelial cells and cholestasis in
768 mice, *Arch. Toxicol.* 94 (2020) 219–229. <https://doi.org/10.1007/s00204-019-02582-8>

4 ARTIGO II

ARTIGO SUBMETIDO PARA PUBLICAÇÃO

**EXPRESSION PROFILE OF KEY GENES INVOLVED IN DNA REPAIR
MECHANISMS IN BOVINE CUMULUS CELLS CULTURED WITH BOVINE
SERUM ALBUMIN OR FETAL CALF SERUM**

André Lucio Fontana Goetten, Júlia Koch, Cecília Constantino Rocha, Alceu Mezzalira,
Christopher Allan Price, Valério Marques Portela, Marcos Henrique Barreta

Expression profile of key genes involved in DNA repair mechanisms in bovine cumulus cells cultured with bovine serum albumin or fetal calf serum

André Lucio Fontana Goetten^a, Júlia Koch^a, Cecília Constantino Rocha^b, Alceu Mezzalira^c, Christopher Allan Price^d, Valério Marques Portela^e & Marcos Henrique Barreta^{a*}

^a Centro de Ciências Rurais, Universidade Federal de Santa Catarina (UFSC) - Curitibanos, SC, Brazil.

^b Department of Animal Sciences, University of Florida, Gainesville, FL, USA.

^c Laboratório de Reprodução Animal Assis Roberto de Bem, Centro de Ciências Agroveterinárias (CAV), Universidade do Estado de Santa Catarina (UDESC) - Lages, SC, Brazil.

^d Centre de Recherche en Reproduction Animale, Faculté de Médecine Vétérinaire, Université de Montréal - St-Hyacinthe, Quebec, Canada.

^e Laboratório de Biotecnologia e Reprodução Animal (BIOREP), Universidade Federal de Santa Maria (UFSM) - Santa Maria, RS, Brazil.

* Corresponding author at: Centro de Ciências Rurais, Universidade Federal de Santa Catarina (UFSC) – Campus Curitibanos, Rodovia Ulysses Gaboardi, km 03, 89.520-000, Curitibanos, SC, Brazil. TEL: +55(48)3721-4731. E-mail address: marcos.barreta@ufsc.br (M.H. Barreta)

Acknowledgements. The authors would like to thank El’Golli and Verdi abattoirs for providing the bovine ovaries.

4.1 ABSTRACT

Cumulus cells from cumulus-oocyte complexes (COC) matured in vitro in serum-free medium show high incidence of apoptosis and DNA double-strand breaks (DSB). This study aimed to characterize the transcript expression profile of selected genes involved in DNA repair mechanisms in bovine cumulus cells cultured with bovine serum albumin (BSA) or fetal calf serum (FCS). Briefly, bovine cumulus-oocyte complexes were in vitro matured with either, 0,4 % BSA or 10% FCS for 3, 6, 12 or 24 hours. The total RNA of cumulus cells was used for real-time PCR analysis. The mRNA expression was regulated during the 24 h of IVM for *XRCC6*, *XRCC5*, *DNAPK*, *GAAD45B*, *TP53BP1*, *RAD50*, *RAD52*, *ATM* and *BRCA2* target genes ($P < 0.05$). Although, an interaction between protein source (FCS or BSA) and time was not detected ($P \geq 0.05$). Cumulus cells from COCs matured with BSA presented higher mRNA expression of two genes compared to FCS group: *TP53BP1* at 6 h and *BRCA1* at 3, 6, 12 and 24 h ($P < 0.05$). In summary, our results showed for the first time the expression profile of the key genes involved in DSB repair mechanisms in cumulus cells obtained from bovine COCs matured with FCS or BSA. The higher mRNA expression of *BRCA1* and *TP53BP1* and lower mRNA expression of *TNFAIP6* suggests an increase in apoptosis rate and DNA damage in cumulus cells cultured in BSA-supplemented medium and may explain, at least to some extent, the reduced developmental potential of bovine oocytes matured in serum-free medium.

Keywords: In vitro maturation, cumulus cells, DNA repair, apoptosis.

4.2 INTRODUCTION

In vitro maturation (IVM) of oocytes is the major approach in assisted reproductive technologies in cattle, humans and other species [1]. It was designed to obtain mature oocytes

following culture of immature cumulus–oocyte complexes (COC) collected from antral follicles [2]. Despite its widespread use, the efficacy of IVM is lower than that observed for in vivo matured oocytes. Therefore, more detailed information on the molecular pathways that control cumulus cell expansion and COC maturation are needed to improve IVM protocols [1,2].

Cumulus cells plays a fundamental role in proper oocyte maturation and acquisition of further developmental competence. Denuded oocytes when submitted to IMV showed lower competence for embryonic development than cumulus-enclosed oocytes [3]. Cumulus-oocyte communication mechanisms are established and involve bidirectional exchanges through gap junctions and paracrine signaling regulating this phenomenon. Gap junctions from COCs allow the exchange of the small molecules, such as cAMP and cGMP that fundamentally regulate the process of oocyte maturation, directing oocyte progression through meiosis. In addition, the oocyte secretes growth-differentiation factor 9 (*GDF9*) and bone morphogenetic protein 15 (*BMP15*), which are paracrine factors that regulate the biological function of cumulus cells and orchestrates many of the associated local signaling events around ovulation, which is the key to establishing oocyte competence to sustain early embryo development [3–5]. However, this communication is under stress in IVM COCs when compared to those in vivo matured [5].

The combined actions of FSH/cAMP and oocyte-secreted factors (*BMP15* and *GDF9*) induce epidermal growth factors receptors (*EGFR*) activation in COCs [6]. Activation of *EGFR* signaling in cumulus cells, together with cAMP priming, triggers oocyte nuclear maturation and acquisition of developmental competence as well as cumulus expansion, which are crucial for ovulation [7]. In cattle, granulosa cells stimulated by LH, and cumulus cells stimulated by FSH, produce epidermal growth factors (*EGFs*)-Like, mainly epiregulin (*EREG*) and amphiregulin (*AREG*). When *EGFs*-Like binds to *EGFR* expressed in granulosa

and cumulus cells, they stimulate their own synthesis, the synthesis of prostaglandin synthetase 2 (*PTGS2*) and prostaglandin E2 (*PGE2*), as well as the expression of steroidogenesis and cumulus expansion genes [8]. *PGE2* also plays a role in the ovulation process by inducing the synthesis of EGF-like proteins in a similar way as LH does [9]. *EGF-like*, *PTGS2*, and *PGE2* work together to increase the production of tumor necrosis factor 6 inducing protein (*TNFAIP6*), hyalurone synthetase 2 (*HAS2*), and pentraxin 3 (*PTX3*), which are responsible for cumulus expansion [7,9].

Culture conditions during in vitro embryo production (IVP) can impact on the developmental potential of the early embryo. The lower quality of these embryos is due to changes in gene expression patterns because of interaction with the adverse environment generated by in vitro conditions [10]. IVM protein supplementation can have profound effects on development rate and overall developmental efficiency, as demonstrated by morula and blastocysts yield [11]. Most IVM procedures use serum to improve blastocyst developmental rate in cattle, but this has become less attractive because it is associated with a number of problematic issues. In addition to scientific disadvantages such as unknown exact composition, seasonal and geographical variability between batches, serious safety concerns for both laboratory staff and animals due to the risk of transmitting diseases potentially present in bovine serum, as well as ethical concerns about fetal distress have been raised with regard to the use of fetal bovine serum [12]. Therefore, the goal of IVP it is to develop optimal conditions for IVM in serum-free media and bovine serum albumin is the most widely used protein source for serum replacement. Unfortunately, the absence of serum in IVM medium results in reduced blastocyst rate [13–15], lower degree of cumulus expansion [15,16] and higher incidence of apoptosis and DNA fragmentation in cumulus cells [17].

To maintain genomic integrity, DNA must be protected from damage induced by environmental agents or generated spontaneously during DNA metabolism [18]. Among the

different types of DNA damage induced by environmental genotoxins, DNA double-strand breaks (DSB) are the most deleterious type of damage and can substantially alter genetic integrity [19]. Repair of DSBs is mainly coordinated by two molecular pathways: homologous recombination (HR) and non-homologous end-joining (NHEJ). Whereas HR is known as the error-free repair pathway because it uses the sister chromatid as a template for repair, NHEJ is referred to as the error-prone pathway since it does not depend on any template [20]. Growth arrest and DNA damage inducible beta (*GADD45B*) and tumor suppressor P53-binding protein 1 (*TP53BP1*) are mediators of DSB repair which display a complex array of physical interactions with other cellular proteins and, in some cases, these systems interfere with the cell cycle and can activate specific checkpoints in order to repair the damage or induce apoptosis when cell protection is not possible [21–23]. COCs in vitro matured in serum-free media have a high rate of apoptosis in cumulus cells and DSB repair mechanisms can be compromised under these cell culture conditions. The presence of mRNA encoding genes implicated in these processes has been demonstrated in human [24], mouse [19], swine [20], and bovine [25,26] oocytes, as well as in human [27], and rat [28] cumulus cells. However, there is no information about the expression of genes involved in HR and NHEJ pathways in bovine cumulus cells.

Clearly, coordinated bilateral interaction between the cumulus cells and the oocyte is essential for the full ability of both cell types to promote early embryogenesis. Thus, the information stored in the cumulus cells may be direct markers for the further developmental fate of the oocyte. Therefore, gene expression patterns in biopsied cumulus cells might actually serve as non-invasive markers to evaluate oocyte maturation. Changes in oocyte maturation conditions are associated with modifications in the expression profile of several key genes in cumulus cells, since bovine cumulus cells isolated from in vitro matured COCs have a different transcriptomic signature than cumulus cells isolated from in vivo matured

COCs [29]. This is why the profile expression of DNA repair genes in bovine cumulus cells is an intriguing question that could provide knowledge to improve the efficiency of IVM. Thus, the aims of this study were to: (1) characterize the transcript expression profile of selected genes involved in DNA DSB repair mechanisms in bovine cumulus cells obtained from COCs in vitro matured; and (2) investigate whether supplementing IVM medium with bovine serum albumin (BSA) or fetal calf serum (FCS) affects transcript expression of these genes.

4.3 MATERIAL AND METHODS

All reagents were purchased from Sigma-Aldrich Co. (Rocklin, CA, USA) unless otherwise stated.

4.3.1 In vitro maturation of Cumulus-oocyte complexes

Cow ovaries were obtained from a local abattoir and transported to the laboratory in saline solution (0.9% NaCl) containing 100 IU/mL penicillin and 50 µg/mL streptomycin sulfate at 25°C. Cumulus oocyte complexes (COCs) from 3 to 8 mm diameter follicles were aspirated with a vacuum pump (vacuum rate of 15mL of water/minute) and pooled in a 15 mL conical tube. After sedimentation, COCs were recovered and selected according to Leibfried & First [30] under a stereomicroscope. Three independent replicates of grade 1 or 2 COCs (n=15/group/replicate), with each replicate using ovaries collected at different days, were randomly transferred to 4-well plates into 400 µL of maturation medium with bovine serum albumin (BSA) or fetal calf serum (FCS) and cultured in an incubator at 38.5°C in a saturated humidity atmosphere containing 5% CO₂ and 95% air. The maturation medium used was TCM199 containing Earle's salts and L-glutamine supplemented with 25 mM HEPES, 0.2 mM pyruvic acid, 2.2 mg/mL sodium bicarbonate, 5.0 µg/mL LH (Lutropin-V, Bioniche Animal Health Canada Inc., Belleville, ON, Canada), 0.1 µg/mL FSH (Folltropin-V, Bioniche Animal Health Canada Inc., Belleville, ON, Canada), 100 IU/mL penicillin, 50 µg/mL

streptomycin sulfate and 0.4% fatty acid-free BSA (BSA group) or 10% FCS (FCS group). COCs were cultured for 3, 6, 12 or 24 hours and cumulus cells were recovered by vortexing, collected into Trizol and stored at -80°C until RNA extraction.

4.3.2 Nucleic acid extraction and qPCR

Total RNA was extracted using Trizol® (Thermo Fisher Scientific Inc. Waltham, MA, USA) according to the manufacturer's instructions. The concentration and purity of total RNA extracts were evaluated using a spectrophotometer (NanoVue, GE Healthcare, Cambridge, England). Total RNA (200 ng) was first treated with gDNA Wipeout Buffer (Qiagen, Mississauga, ON, Canada) at 42°C for 2 minutes to digest any contaminating DNA. Then RNA was reverse transcribed (RT) in the presence of Quantiscript Reverse Transcriptase and RT primer mix (QuantiTec RT Kit® - Qiagen, Mississauga, ON, Canada) in a volume of 20 µL at 42°C for 15 minutes. The reaction was stopped by incubation at 95°C for 3 minutes. The cDNA was stored at -20°C until quantitative PCR (qPCR) analysis. Quantification of specific transcripts was performed by RT-qPCR using QuantiNova SYBR Green PCR Kit® (Qiagen, Mississauga, ON, Canada), and reactions were carried out using a Stratagene MX3005P apparatus (Agilent, Santa Clara, CA, USA). Common thermal cycling parameters (3 minutes at 95°C, 40 cycles of 15 seconds at 95°C, 30 seconds at 60°C, and 30 seconds at 72°C) were used to amplify each transcript. The reaction was performed in duplicate, and melting-curves analyzes were performed to verify product identity. The selected target genes were epiregulin (*EREG*), hyaluronan synthase 2 (*HAS2*), prostaglandin-endoperoxide synthase 2 (*PTGS2*), TNF alpha induced protein 6 (*TNFAIP6*), growth arrest and DNA damage inducible beta (*GADD45B*), tumor protein p53 binding protein 1 (*TP53BP1*), X-ray repair cross complementing 5 (*XRCC5*), X-ray repair cross complementing 6 (*XRCC6*), Protein kinase DNA-activated, catalytic subunit (*DNAPK*), ATM serine/threonine kinase (*ATM*), BRCA1 DNA repair associated (*BRCA1*), BRCA2 DNA repair associated (*BRCA2*),

RAD50 double-strand break repair protein (*RAD50*), RAD51 recombinase (*RAD51*) and RAD52 homolog DNA repair protein (*RAD52*). The genes H2A histone family, member Z (*H2AFZ*), cyclophilin B (*PPIB*), and TATA-box binding protein (*TBP*) were tested as reference genes. The primers (Table 1) were previously described for cattle. The amplification data were extracted from the Stratagene MX3005P apparatus, and each sample was analyzed through LinReg PCR software [31] for baseline correction and determination of qPCR efficiency. To select reference genes, geNorm software [32] was used. *PPIB* and *TPB* were selected as the best reference genes. The expression of each target gene relative to the expression of the reference genes was calculated using the Pfaffl method with efficiency correction and normalized to a calibrator sample [33].

4.3.3 Statistical analysis

All statistical analyzes were performed using the R 4.1.1 software (R Core Team, 2021, Vienna, Austria). For the mRNA expression, the assumption of normality was checked by Shapiro-Wilk test and homogeneity of variance was tested by Bartlett test. Data that did not follow normal distribution according to the Shapiro–Wilk test, or either, exhibit homoscedasticity according to the Bartlett test were transformed by Box-Cox power transformation. Thereafter, the data were analyzed using a linear model by an ANOVA for two-way factorial design (R Package “ExpDes.pt” version 1.2.1). The main effects of protein source (BSA and FCS), time (3, 6, 12, and 24 h of IVM) and the interaction were tested. In case of significant effects in the model ($p < 0.05$), comparisons were performed using the least-square means (LSMeans) method (R Package “emmeans” version 1.6.3) with Tukey-Kramer HSD test for multiple comparisons adjustment. All data are presented as means $\pm$ standard error of mean (S.E.M.) of three independent replicates.

4.4 RESULTS

The relative expression of *PTGS2*, *EREG*, *HAS2* and *TNFAIP6* mRNA was affected over the 24 h of IVM ($P < 0.05$) (Figure 1). Only maturation time effect was found for the expression of *PTGS2*, *EREG* and *HAS2*, regardless of protein source supplied. The relative abundance of *PTGS2* mRNA increased gradually up to 24 h of IVM (Figure 1A). A significant increase in *EREG* transcript abundance was detected at 6 h and 12 h of IVM as compared to the beginning (3 h) and the end (24 h) of IVM (Figure 1B). *HAS2* expression was characterized by a temporary increase at 6 h of IVM, followed by a significant decrease at 12 h of IVM and then remained relatively stable up to 24 h of IVM (Figure 1C). When evaluating *TNFAIP6* expression, significant effects of protein source, time and their interaction were detected (Figure 1D). The protein source effect indicated that *TNFAIP6* expression was 1.6-fold greater in cumulus cells cultured in FCS-supplemented medium than in those cultured in BSA-supplemented medium. The interaction between protein source and time was demonstrated by a marked increase in *TNFAIP6* expression in the FCS group between 12 and 24 h of IVM. Whereas the most significant increase in *TNFAIP6* transcript in the BSA group occurred only at 24 h of IVM. In addition, the FCS group had more *TNFAIP6* transcripts than the BSA group at 12 and 24 h of IVM.

As shown in Figure 2, although a significant ($P < 0.05$) effect of maturation time was detected, there was no interaction between protein source and time in *XRCC6*, *XRCC5*, *DNAPK*, *GADD45B* and *TP53BP1* mRNA expression. For *TP53BP1*, significant effects of protein source and time were detected (Figure 2A). There was a 1.33-fold higher level of *TP53BP1* expression in cumulus cells cultivated in BSA-supplemented medium compared to cumulus cells cultured in FCS-supplemented medium, indicating a protein source effect. The time effect was demonstrated by relatively stable *TP53BP1* expression between 3 and 6 h of IVM, followed by a progressive increase from 12 h of IVM, regardless of protein source

supplemented. Furthermore, cells supplemented with BSA showed higher expression of TP53BP1 than cells supplemented with FCS at 6 h of IVM. The abundance of *XRCC6* transcripts (Figure 2B) showed a significant increase only at 24 h of IVM, with no changes throughout the remainder of the culture period, regardless of the use of BSA or FCS in IVM. After a sudden drop at 6 h of IVM, *XRCC5* expression (Figure 2C) increased progressively and significantly until 24 h of IVM, irrespective of the protein source in the culture medium. *DNAPK* and *GADD45* exhibited almost the same mRNA expression pattern (Figures 2D and 2E). The abundance of both *DNAPK* and *GADD45* transcripts remained relatively unchanged until 6 h of IVM, increased significantly at 12 h of IVM and then maintained relatively constant until 24 h of IVM.

Expression of the six HR pathway mediators investigated is shown in Figure 3. *RAD50*, *RAD52*, *ATM* and *BRCA2* were expressed similarly over time (Figures 3A, 3C, 3D and 3F). After a decrease at 6 h of IVM, the expression levels significantly increased up to 24 h of IVM, indicating a time effect ($P < 0.05$) for these four genes. A protein source effect ($P < 0.05$) was found for *BRCA1*, as evidenced by a 1.45-fold higher expression of *BRCA1* in cumulus cells cultivated with BSA versus FCS-supplemented (Figure 3E), but there was no significant variation in *BRCA1* transcript abundance at different time points ($P > 0.05$). The abundance of *RAD51* transcripts remained relatively constant over the course of the IVM ($P > 0.05$), regardless of the protein source supplemented (Figure 3B). There was no interaction between protein source and time for any of the HR mediators studied ($P > 0.05$).

4.5 DISCUSSION

In this study, COCs were cultured in order to characterize the transcript expression profile of selected genes involved in DNA DSB repair mechanisms in bovine cumulus cells matured in vitro and to investigate whether supplementing IVM medium with BSA or FCS affects the abundance of these transcripts. COCs were collected at four separate IVM time

points, corresponding to the different stages of maturation progression [34]: 3 hours - germinal vesicle; 6 hours - germinal vesicle breakdown; 12 hours - metaphase-I (because oocytes with DNA damage are arrested at this stage [35]; 24 hours - metaphase-II. Furthermore, each of these moments is associated with the in vitro expression of key genes of the ovulatory cascade that are responsible for cumulus expansion. Thus, increased abundance of *HAS2* is expected between 4 and 8 hours of IVM; *EREG* until 12 hours of IVM; and *PTGS2* and *TNFAIP6* starting at 12 hours of IVM [1,36,37]. Since the expression of *EREG*, *HAS2*, *PTGS2* and *TNFAIP6* mRNA in cumulus cells was associated with an increase in oocyte developmental competence and embryo quality [29,38], the levels of mRNA encoding these typical genes upregulated during the preovulatory cascade were assessed to confirm that our IVM system was functional. In most mammals, the EGF signaling network regulates the expression of key genes involved in gap junction closure and production of a large extracellular matrix by cumulus cells, as well as the downregulation of the meiotic inhibitory signaling network, which leads to cumulus expansion and oocyte meiotic maturation [9]. In this study, transient upregulation of the *EREG* transcript was followed by brief *HAS2* upregulation and persistent *PTGS2* and *TNFAIP6* upregulation, as expected. The mRNA expression patterns of these genes over time were consistent with those previously published in cattle COCs [1]. Therefore, it confirms that COCs responded to the IVM protocol and that the dynamic bidirectional COC communication was preserved.

Repair of DSBs involves two pathways, homologous recombination (HR) or non-homologous end-joining (NHEJ). The HR commonly occurs during the S and G2-M phases of the cell cycle [21,39]. A complex of proteins including RAD50, NSB1 and ATM generate single-stranded overhangs at the point of DNA damage, which are then used to match homologous sequences in the sister chromatid. The DNA repair is then affected by multiple mediator proteins including RAD51, RAD52, BRCA1 and BRCA2 [40]. The tumor

suppressor protein BRCA1 is a crucial member of the ATM-mediated DSB repair family. It plays an important role in maintaining genetic integrity via interaction with a number of other proteins, including MRE11, RAD50 and BRCA2 [41,42].

During NHEJ, Ku70 and Ku80 (products of the *XRCC6* and *XRCC5* genes, respectively) are pivotal proteins involved in DNA damage repair. Ku70 cooperates with Ku80 to form Ku heterodimer, which initiates NHEJ pathways in DSBs repair. Ku heterodimer bind to DNA ends and recruit the *DNAPK*. Once bound, *DNAPK* activates its own catalytic subunit (*DNAPKcs*) and further enlists the endonuclease *Artemis*. At a subset of DSBs, *Artemis* removes excess single-strand DNA and generates a substrate that will be ligated by *DNA-ligase-IV*. NHEJ involves blunt-end ligation independent of sequence homology. This pathway can occur throughout the cell cycle but is dominant in G0/G1 and G2 stage of the cell cycle [20,43]. Recently, Ku70 have been proved crucial in some fundamental cellular processes, metabolism, ageing, and related diseases, such as premature ovarian insufficiency, cancers, diabetes, neurodegenerative and cardiovascular diseases [44–47]. In hepatocarcinoma cell-lines, reduction in mRNA of these three genes coding the NHEJ-initiating proteins was related to the stress caused by serum starvation compared to serum-fed cells [48]. In this work, the temporal expression pattern of the DSB repair genes followed a similar trend across time: a decrease or steady abundance of transcripts between 3 and 6 h of IVM, followed by an increase from 6 h onwards. The highest relative abundance of RNA was observed at 24 hours of IVM. The *BRCA1* expression was an exception, since it did not change over the course of maturation.

Regarding the source of protein added to IVM medium, differences were found in the expression of only three genes: *TNFAIP6*, *BRCA1* and *TP53BP1*.

Optimal expansion of the cumulus mass seems to be important for ovulation in mice [49] and this idea was supported by an experiment in which targeted disruption of *TNFAIP6*

[50] led to severe infertility due to impaired cumulus expansion and a defect in its organization. The presence of *TNFAIP6* protein in porcine COCs is involved in stabilizing the mucoelastic extracellular matrix [38]. In cattle abundance of *TNFAIP6* mRNA was significantly higher in cumulus cells from competent oocytes compared to incompetent oocytes [51], and in cumulus cells derived from in vivo COCs compared with their in vitro counterparts [29].

In the present study, *TNFAIP6* expression increased from 12 h of IVM in cumulus cells cultured in FCS-supplemented medium and only at 24 h of IVM in BSA-supplemented medium, with greater transcript abundance in the FCS group than in the BSA group at 12 h and 24 h of IVM. These results are consistent with those reported in a previous study that showed a higher degree of expansion in COCs submitted to IVM in medium supplemented with FCS, compared with those submitted to IVM in medium supplemented with BSA [16], and could be attributed to FCS components that act as transcription factors for genes involved in both extracellular matrix expansion and mucification.

BRCA1 participates in several cellular processes in response to DNA damage. The expression of *BRCA1* and other genes related to the *ATM* repair pathway was significantly increased in the aging and apoptotic cumulus cells of aging women [52]. Our data showed for the first time that cumulus cells cultivated with BSA have higher expression of *BRCA1* and *TP53BP1* than those cultivated with FCS. *TP53BP1* is a key regulator of DSB repair [53]. It functions at the intersection of two major DSB repair pathways (HR and NHEJ) and integrates cellular inputs to ensure the timely execution of repair in the proper cellular contexts [22]. DNA damage can be induced by endogenous metabolites that can react with macromolecules, such as lipids, proteins or nucleic acids and exogenous stimulants, such as ionizing radiation, ultraviolet radiation and reactive oxygen species (ROS) [57–60]. An increase in *TP53BP1* and *BRCA1* mRNA expression was observed after the induction of ROS

production in cumulus cells obtained from vitrified oocytes [54]. Although the presence of cumulus cells during in vitro fertilization protects the bovine oocyte against oxidative stress and improves the first cleavage [55], bovine COCs matured in serum-free medium produce lower progesterone, with higher rate of apoptosis and a lower capacity for embryonic development compared with their FCS-supplemented counterparts [56]. Therefore, higher levels of *BRCA1* and *TP53BP1* mRNA in cumulus cells cultivated in serum-free medium may be related with elevated rates of apoptosis. As a result, it impairs the competence of COCs for embryonic development.

In our IVM model, the HR mediator proteins *RAD50*, *RAD51*, *RAD52*, *ATM* and *BRCA2* were not differentially expressed in cumulus cells cultured with BSA or FCS and although serum suppression in culture systems has potential to reduce the expression of NHEJ-initiating proteins, this find was not observed.

In summary, our results showed for the first time the profile expression of key genes involved in the mechanism of double-strand DNA repair in bovine cumulus cells of COCs matured with FCS or BSA. The higher mRNA expression of *BRCA1* and *TP53BP1* and lower mRNA expression of *TNFAIP6* suggests an increase in apoptosis rate and DNA damage in cumulus cells cultured in BSA-supplemented medium and may explain, at least to some extent, the reduced developmental potential of bovine oocytes matured in serum-free medium.

Source of Funding

This project was supported by the Brazilian National Council for Scientific and Technological Development (CNPq) and Santa Catarina State Research and Innovation Support Foundation (FAPESC; TR2012000474).

Conflict of interest

The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

347

348 4.6 REFERENCES

- 349 [1] Caixeta ES, Sutton-McDowall ML, Gilchrist RB, Thompson JG, Price CA, Machado
 350 MF, et al. Bone morphogenetic protein 15 and fibroblast growth factor 10 enhance
 351 cumulus expansion, glucose uptake, and expression of genes in the ovulatory cascade
 352 during in vitro maturation of bovine cumulus-oocyte complexes. *Reproduction*
 353 2013;146:27–35. <https://doi.org/10.1530/REP-13-0079>.
- 354 [2] De Vos M, Grynberg M, Ho TM, Yuan Y, Albertini DF, Gilchrist RB. Perspectives on
 355 the development and future of oocyte IVM in clinical practice. *J Assist Reprod Genet*
 356 2021;38:1265–80. <https://doi.org/10.1007/s10815-021-02263-5>.
- 357 [3] Tanghe S, Van Soom A, Nauwynck H, Coryn M, de Kruif A. Minireview: Functions of
 358 the cumulus oophorus during oocyte maturation, ovulation, and fertilization. *Mol*
 359 *Reprod Dev* 2002;61:414–24.
- 360 [4] Richani D, Dunning KR, Thompson JG, Gilchrist RB. Metabolic co-dependence of the
 361 oocyte and cumulus cells: essential role in determining oocyte developmental
 362 competence. *Hum Reprod Update* 2021;27:27–47.
 363 <https://doi.org/10.1093/humupd/dmaa043>.
- 364 [5] Brown HM, Dunning KR, Sutton-McDowall M, Gilchrist RB, Thompson JG, Russell
 365 DL. Failure to launch: aberrant cumulus gene expression during oocyte in vitro
 366 maturation. *Reproduction* 2017;153:R109–20. <https://doi.org/10.1530/REP-16-0426>.
- 367 [6] Sugimura S, Richani D, Gilchrist RB. Follicular guidance for oocyte developmental
 368 competence. *Anim Reprod* 2018;15:721–6. [https://doi.org/10.21451/1984-3143-](https://doi.org/10.21451/1984-3143-AR2018-0035)
 369 [AR2018-0035](https://doi.org/10.21451/1984-3143-AR2018-0035).
- 370 [7] Conti M, Hsieh M, Park J-Y, Su Y-Q. Role of the Epidermal Growth Factor Network
 371 in Ovarian Follicles. *Mol Endocrinol* 2006;20:715–23.

- 372 <https://doi.org/10.1210/me.2005-0185>.
- 373 [8] Ben-Ami I, Freimann S, Armon L, Dantes A, Ron-El R, Amsterdam A. Novel function
374 of ovarian growth factors: Combined studies by DNA microarray, biochemical and
375 physiological approaches. *Mol Hum Reprod* 2006;12:413–9.
376 <https://doi.org/10.1093/molehr/gal045>.
- 377 [9] Shimada M, Hernandez-Gonzalez I, Gonzalez-Robayna I, Richards JS. Paracrine and
378 Autocrine Regulation of Epidermal Growth Factor-Like Factors in Cumulus Oocyte
379 Complexes and Granulosa Cells: Key Roles for Prostaglandin Synthase 2 and
380 Progesterone Receptor. *Mol Endocrinol* 2006;20:1352–65.
381 <https://doi.org/10.1210/me.2005-0504>.
- 382 [10] Lonergan P, Rizos D, Gutierrez-Adan A, Fair T, Boland M. Oocyte and Embryo
383 Quality: Effect of Origin, Culture Conditions and Gene Expression Patterns. *Reprod*
384 *Domest Anim* 2003;38:259–67. <https://doi.org/10.1046/j.1439-0531.2003.00437.x>.
- 385 [11] Ali A, Sirard M-A. Effect of the Absence or Presence of Various Protein Supplements
386 on Further Development of Bovine Oocytes During In Vitro Maturation¹. *Biol Reprod*
387 2002;66:901–5. <https://doi.org/10.1095/biolreprod66.4.901>.
- 388 [12] van der Valk J, Bieback K, Buta C, Cochrane B, Dirks WG, Fu J, et al. Fetal bovine
389 serum (FBS): Past – present – future. *ALTEX* 2018;35:99–118.
390 <https://doi.org/10.14573/altex.1705101>.
- 391 [13] Rizos D, Gutiérrez-Adán A, Pérez-Garnelo S, de la Fuente J, Boland MP, Lonergan P.
392 Bovine Embryo Culture in the Presence or Absence of Serum: Implications for
393 Blastocyst Development, Cryotolerance, and Messenger RNA Expression. *Biol Reprod*
394 2003;68:236–43. <https://doi.org/10.1095/biolreprod.102.007799>.
- 395 [14] del Collado M, Saraiva NZ, Lopes FL, Gaspar RC, Padilha LC, Costa RR, et al.
396 Influence of bovine serum albumin and fetal bovine serum supplementation during in

- 397 vitro maturation on lipid and mitochondrial behaviour in oocytes and lipid
 398 accumulation in bovine embryos. *Reprod Fertil Dev* 2015;28:1721–32.
 399 <https://doi.org/10.1071/RD15067>.
- 400 [15] Rätty M, Ketoja E, Pitkänen T, Ahola V, Kananen K, Peippo J. In vitro maturation
 401 supplements affect developmental competence of bovine cumulus–oocyte complexes
 402 and embryo quality after vitrification. *Cryobiology* 2011;63:245–55.
 403 <https://doi.org/10.1016/j.cryobiol.2011.09.134>.
- 404 [16] Velho FA de B, Costa BG, Alcoba DD, Braga BL da R, Monroy NLS, Oliveira ATD,
 405 et al. Gene Expression and In Vitro Nuclear Maturation in Bovine Cumulus Oocyte
 406 Complexes Matured in a Medium Supplemented with Bovine Fetal Serum or Bovine
 407 Serum Albumin. *Acta Sci Vet* 2013;41:01–7.
- 408 [17] Ikeda S, Imai H, Yamada M. Apoptosis in cumulus cells during in vitro maturation of
 409 bovine cumulus-enclosed oocytes. *Reproduction* 2003;125:369–76.
 410 <https://doi.org/10.1530/rep.0.1250369>.
- 411 [18] Ciccia A, Elledge SJ. The DNA Damage Response: Making It Safe to Play with
 412 Knives. *Mol Cell* 2010;40:179–204. <https://doi.org/10.1016/j.molcel.2010.09.019>.
- 413 [19] Titus S, Li F, Stobezki R, Akula K, Unsal E, Jeong K, et al. Impairment of BRCA1-
 414 Related DNA Double-Strand Break Repair Leads to Ovarian Aging in Mice and
 415 Humans. *Sci Transl Med* 2013;5. <https://doi.org/10.1126/scitranslmed.3004925>.
- 416 [20] Bohrer RC, Dicks N, Gutierrez K, Duggavathi R, Bordignon V. Double-strand DNA
 417 breaks are mainly repaired by the homologous recombination pathway in early
 418 developing swine embryos. *FASEB J* 2018;32:1818–29.
 419 <https://doi.org/10.1096/fj.201700800R>.
- 420 [21] Lieber MR, Gu J, Lu H, Shimazaki N, Tsai AG. Nonhomologous DNA end joining
 421 (NHEJ) and chromosomal translocations in humans. *Subcell Biochem* 2010;50:279–96.

- 422 [22] Panier S, Boulton SJ. Double-strand break repair: 53BP1 comes into focus. *Nat Rev*
 423 *Mol Cell Biol* 2014;15:7–18. <https://doi.org/10.1038/nrm3719>.
- 424 [23] Hoffman B, Liebermann DA. Role of gadd45 in myeloid cells in response to
 425 hematopoietic stress. *Blood Cells, Mol Dis* 2007;39:344–7.
 426 <https://doi.org/10.1016/j.bcmed.2007.06.011>.
- 427 [24] Jaroudi S, Kakourou G, Cawood S, Doshi A, Ranieri DM, Serhal P, et al. Expression
 428 profiling of DNA repair genes in human oocytes and blastocysts using microarrays.
 429 *Hum Reprod* 2009;24:2649–55. <https://doi.org/10.1093/humrep/dep224>.
- 430 [25] Bilotto S, Boni R, Russo GL, Lioi MB. Meiosis progression and donor age affect
 431 expression profile of DNA repair genes in bovine oocytes. *Zygote* 2015;23:11–8.
 432 <https://doi.org/10.1017/S0967199413000154>.
- 433 [26] Barreta MH, Gasperin BG, Rissi VB, Cesaro MP de, Ferreira R, de Oliveira JF, et al.
 434 Homologous recombination and non-homologous end-joining repair pathways in
 435 bovine embryos with different developmental competence. *Exp Cell Res*
 436 2012;318:2049–58. <https://doi.org/10.1016/j.yexcr.2012.06.003>.
- 437 [27] Ouandaogo ZG, Frydman N, Hesters L, Assou S, Haouzi D, Dechaud H, et al.
 438 Differences in transcriptomic profiles of human cumulus cells isolated from oocytes at
 439 GV, MI and MII stages after in vivo and in vitro oocyte maturation. *Hum Reprod*
 440 2012;27:2438–47. <https://doi.org/10.1093/humrep/des172>.
- 441 [28] Agca C, Yakan A, Agca Y. Estrus synchronization and ovarian hyper-stimulation
 442 treatments have negligible effects on cumulus oocyte complex gene expression
 443 whereas induction of ovulation causes major expression changes. *Mol Reprod Dev*
 444 2013;80:102–17. <https://doi.org/10.1002/mrd.22141>.
- 445 [29] Tesfaye D, Ghanem N, Carter F, Fair T, Sirard M-A, Hoelker M, et al. Gene expression
 446 profile of cumulus cells derived from cumulus - oocyte complexes matured either in

- 447 vivo or in vitro. *Reprod Fertil Dev* 2009;21:451. <https://doi.org/10.1071/RD08190>.
- 448 [30] Leibfried L, First NL. Characterization of bovine follicular oocytes and their ability to
449 mature. *J Anim Sci* 1979;48:76–86. <https://doi.org/10.2527/jas1979.48176x>.
- 450 [31] Ramakers C, Ruijter JM, Lekanane RH, Moorman AFM. Assumption-free analysis of
451 quantitative real-time polymerase chain reaction (PCR) data 2003;339:62–6.
452 [https://doi.org/10.1016/S0304-3940\(02\)01423-4](https://doi.org/10.1016/S0304-3940(02)01423-4).
- 453 [32] Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, et al.
454 Accurate normalization of real-time quantitative RT-PCR data by geometric averaging
455 of multiple internal control genes. *Genome Biol* 2002;3:1–12.
456 <https://doi.org/https://doi.org/10.1186/gb-2002-3-7-research0034>.
- 457 [33] Pfaffl MW. A new mathematical model for relative quantification in real-time RT-
458 PCR. *Nucleic Acids Res* 2001;29:2002–7. <https://doi.org/10.1093/nar/29.9.e45>.
- 459 [34] Sirard MA, Florman HM, Leibfried-Rutledge ML, Barnes FL, Sims ML, First NL.
460 Timing of Nuclear Progression and Protein Synthesis Necessary for Meiotic
461 Maturation of Bovine Oocytes. *Biol Reprod* 1989;40:1257–63.
462 <https://doi.org/10.1095/biolreprod40.6.1257>.
- 463 [35] Marangos P, Stevense M, Niaka K, Lagoudaki M, Nabti I, Jessberger R, et al. DNA
464 damage-induced metaphase I arrest is mediated by the spindle assembly checkpoint and
465 maternal age. *Nat Commun* 2015;6:8706. <https://doi.org/10.1038/ncomms9706>.
- 466 [36] Caixeta ES, Machado MF, Ripamonte P, Price C, Buratini J. Effects of FSH on the
467 expression of receptors for oocyte-secreted factors and members of the EGF-like
468 family during in vitro maturation in cattle. *Reprod Fertil Dev* 2013;25:890–9.
469 <https://doi.org/10.1071/RD12125>.
- 470 [37] Machado MF, Caixeta ES, Sudiman J, Gilchrist RB, Thompson JG, Lima PF, et al.
471 Fibroblast growth factor 17 and bone morphogenetic protein 15 enhance cumulus

- expansion and improve quality of in vitro-produced embryos in cattle. *Theriogenology* 2015;84:390–8. <https://doi.org/10.1016/j.theriogenology.2015.03.031>.
- [38] Nagyova E, Camaioni A, Prochazka R, Day AJ, Salustri A. Synthesis of Tumor Necrosis Factor Alpha-Induced Protein 6 in Porcine Preovulatory Follicles : A Study with A38 Antibody. *Biol Reprod* 2008;78:903–9. <https://doi.org/10.1095/biolreprod.107.064832>.
- [39] Feng L, Li N, Li Y, Wang J, Gao M, Wang W, et al. Cell cycle-dependent inhibition of 53BP1 signaling by BRCA1. *Cell Discov* 2015;1:15019. <https://doi.org/10.1038/celldisc.2015.19>.
- [40] Ceccaldi R, Rondinelli B, D’Andrea AD. Repair Pathway Choices and Consequences at the Double-Strand Break. *Trends Cell Biol* 2016;26:52–64. <https://doi.org/10.1016/j.tcb.2015.07.009>.
- [41] Gudmundsdottir K, Ashworth A. The roles of BRCA1 and BRCA2 and associated proteins in the maintenance of genomic stability. *Oncogene* 2006;25:5864–74. <https://doi.org/10.1038/sj.onc.1209874>.
- [42] Venkitaraman AR. Cancer Susceptibility and the Functions of BRCA1 and BRCA2. *Cell* 2002;108:171–82. [https://doi.org/10.1016/S0092-8674\(02\)00615-3](https://doi.org/10.1016/S0092-8674(02)00615-3).
- [43] Downs JA, Jackson SP. A means to a DNA end: the many roles of Ku. *Nat Rev Mol Cell Biol* 2004;5:367–78. <https://doi.org/10.1038/nrm1367>.
- [44] Bai P, Cantó C. The Role of PARP-1 and PARP-2 Enzymes in Metabolic Regulation and Disease. *Cell Metab* 2012;16:290–5. <https://doi.org/10.1016/j.cmet.2012.06.016>.
- [45] Bürkle A, Beneke S, Muir ML. Poly(ADP-ribosyl)ation and aging. *Exp Gerontol* 2004;39:1599–601. <https://doi.org/10.1016/j.exger.2004.07.010>.
- [46] Shimizu I, Yoshida Y, Suda M, Minamino T. DNA Damage Response and Metabolic Disease. *Cell Metab* 2014;20:967–77. <https://doi.org/10.1016/j.cmet.2014.10.008>.

- 497 [47] Dang Y, Wang X, Hao Y, Zhang X, Zhao S, Ma J, et al. MicroRNA-379-5p is
 498 associated with biochemical premature ovarian insufficiency through PARP1 and
 499 XRCC6. *Cell Death Dis* 2018;9:106. <https://doi.org/10.1038/s41419-017-0163-8>.
- 500 [48] Zhang T, Zhang X, Shi W, Xu J, Fan H, Zhang S, et al. The DNA damage repair
 501 protein Ku70 regulates tumor cell and hepatic carcinogenesis by interacting with
 502 FOXO4. *Pathol - Res Pract* 2016;212:153–61.
 503 <https://doi.org/10.1016/j.prp.2015.12.012>.
- 504 [49] Chen L, Russell PT, Larsen WJ. Functional Significance of Cumulus Expansion in the
 505 Mouse : Roles for the Preovulatory Synthesis of Hyaluronic Acid Within the Cumulus
 506 Mass. *Mol Reprod Dev* 1993;34:87–93. <https://doi.org/10.1002/mrd.1080340114>.
- 507 [50] Fülöp C, Szántó S, Mukhopadhyay D, Bárdos T, Kamath R V, Rugg MS, et al.
 508 Impaired cumulus mucification and female sterility in tumor necrosis factor-induced
 509 protein-6 deficient mice. *Dev Dis* 2003;130:2253–61.
 510 <https://doi.org/10.1242/dev.00422>.
- 511 [51] Matoba S, Bender K, Fahey AG, Mamo S, Brennan L, Lonergan P, et al. Predictive
 512 value of bovine follicular components as markers of oocyte developmental potential.
 513 *Reprod Fertil Dev* 2014;26:337. <https://doi.org/10.1071/RD13007>.
- 514 [52] Sun X, Jiang H, Han D, Fu Y, Liu J, Gao Y, et al. The activated DNA double-strand
 515 break repair pathway in cumulus cells from aging patients may be used as a convincing
 516 predictor of poor outcomes after in vitro fertilization-embryo transfer treatment. *PLoS*
 517 *One* 2018;13:e0204524. <https://doi.org/10.1371/journal.pone.0204524>.
- 518 [53] Huyen Y, Zgheib O, DiTullio Jr RA, Gorgoulis VG, Zacharatos P, Petty TJ, et al.
 519 Methylated lysine 79 of histone H3 targets 53BP1 to DNA double-strand breaks.
 520 *Nature* 2004;432:406–11. <https://doi.org/10.1038/nature03114>.
- 521 [54] Chang H, Chen H, Zhang L, Wang Y, Xie X, Zhang Y, et al. Effect of oocyte

522 vitrification on DNA damage in metaphase II oocytes and the resulting preimplantation
 523 embryos. *Mol Reprod Dev* 2019;86:1603–14. <https://doi.org/10.1002/mrd.23247>.

- 524 [55] Fatehi AN, Roelen BAJ, Colenbrander B, Schoevers EJ, Gadella BM, Bevers MM, et
 525 al. Presence of cumulus cells during in vitro fertilization protects the bovine oocyte
 526 against oxidative stress and improves first cleavage but does not affect further
 527 development. *Zygote* 2005;13:177–85. <https://doi.org/10.1017/S0967199405003126>.
- 528 [56] Salhab M, Tosca L, Cabau C, Papillier P, Perreau C, Dupont J, et al. Kinetics of gene
 529 expression and signaling in bovine cumulus cells throughout IVM in different mediums
 530 in relation to oocyte developmental competence, cumulus apoptosis and progesterone
 531 secretion. *Theriogenology* 2011;75:90–104.
 532 <https://doi.org/10.1016/j.theriogenology.2010.07.014>.

- 533 [57] Babazadeh B, Sadeghnia HR, Safarpour Kapurchal E, Parsaee H, Nasri S, Tayarani-
 534 Najaran Z. Protective effect of *Nigella sativa* and thymoquinone on serum/glucose
 535 deprivation-induced DNA damage in PC12 cells. *Avicenna J Phytomedicine*
 536 2012;2:125–32.

- 537 [58] Joo J, Hong I, Kim N, Choi E. *Trichosanthes kirilowii* extract enhances repair of UVB
 538 radiation-induced DNA damage by regulating BMAL1 and miR-142-3p in human
 539 keratinocytes. *Mol Med Rep* 2018;17:877–833.
 540 <https://doi.org/10.3892/mmr.2017.7932>.

- 541 [59] Kimeswenger S, Schwarz A, Födinger D, Müller S, Pehamberger H, Schwarz T, et al.
 542 Infrared A radiation promotes survival of human melanocytes carrying ultraviolet
 543 radiation-induced DNA damage. *Exp Dermatol* 2016;25:447–52.
 544 <https://doi.org/10.1111/exd.12968>.

- 545 [60] Salmon TB, Evert BA, Song B, Doetsch PW. Biological consequences of oxidative
 546 stress-induced DNA damage in *Saccharomyces cerevisiae*. *Nucleic Acids Res*

547 2004;32:3712–23. <https://doi.org/10.1093/nar/gkh696>.


548

549

550 **Table 1 – Information of specific primers used for amplification in real-time PCR.**


Gene Symbol	Primer sequence (5' to 3')	Product Size (bp)	Efficiency	Accession Number
<i>Reference genes</i>				
<i>H2AFZ</i>	F: GAGGAGCTGAACAAGCTGTTG R: TTGTGGTGGCTCTCAGTCTTC	104	2,05	XM_002686087.4
<i>PPIB</i>	F: GGTCATCGGTCTCTTTGGAA R: TCCTTGATCACACGATGGAA	117	2,05	NM_174152.2
<i>TBP</i>	F: CAGAGAGCTCCGGGATCGT R: CACCATCTTCCCAGAACTGAATAT	194	2,11	NM_001075742.1
<i>Ovulatory cascade genes needed for cumulus expansion</i>				
<i>EREG</i>	F: CTGCACAGCATTAGTTCAAACCTGA R: TGTCCATGCAAACAGTAGCCATT	100	1,89	XM_010806226.3
<i>HAS2</i>	F: ACACAGACAGGCTGAGGACAACCTT R: AAGCAGCTGTGATTCCAAGGAGGA	133	2,03	NM_174079.3
<i>PTGS2</i>	F: TGCTGAGTTTAAACACGCTCTACCA R: TGAGACCATGTTCCAGTAAGACAGA	125	1,93	NM_174445.2
<i>TNFAIP6</i>	F: GCAAAGGAGTGTGGTGGTGTGTTT R: ACTGAGGTGAATGCGCTGACCATA	135	2,04	NM_001007813.2
<i>Cell cycle checkpoint regulator</i>				
<i>GADD45B</i>	F: TACGAGTCGGCCAAGCTGAT R: GTCCTCCTCTTCCTCGTCGAT	81	2,06	NM_001040604.1
<i>TP53BP1</i>	F: ATCAGACCAACAGCAGAATTTCC R: CACCACGTCAAACACCCCTAA	130	2,01	NM_001206397.2
<i>Non-homologous end-joining repair</i>				
<i>XRCC5</i>	F: TGGCATCTCCCTGCAGTTCT R: AGGCCCATGGTGGTCTGA	100	2,04	NM_001102141.1
<i>XRCC6</i>	F: AATTGACTCCTTTTGACATGAGCAT R: CCATAGAACACCACTGCCAAGA	100	2,00	NM_001192246.1
<i>DNAPK</i>	F: AAAGGCAATCCGTCCTCAGA R: AAGGCAGGTGCTAAACTGAGATG	100	2,05	NM_001256559.2
<i>Homologous recombination repair</i>				
<i>ATM</i>	F: CTTAGGAGGAGCTTGGGCCT R: CCGCTGTGTGGCAAACC	149	2,08	NM_001205935.1
<i>BRCA1</i>	F: ACAAAGCAGCAGACACAATCTCA R: TCATGGTCTCCCACACTGAAATA	110	1,87	NM_178573.1
<i>BRCA2</i>	F: AAATTTCACTGCACCTGGTCAA R: TCATGGGTTTGCCTATAGTTATCG	170	1,77	XM_002691807.5
<i>RAD50</i>	F: TGTGGAACAGGGCCGTCTA R: CAATTCTAGCTGTGTTGCCAGAGA	100	2,02	NM_001206868.1
<i>RAD51</i>	F: ATGCACCGAAGAAGGAGCTAAT R: GATCGCCTTTGGTGGAACTC	130	2,05	NM_001046179.2
<i>RAD52</i>	F: GGCCAGGAAGGAGGCAGTA R: TGACCTCAGATAGTCTTTGTCCAGAA	100	1,98	NM_001024525.1

Fig. 1 – Abundance of A) Prostaglandin-endoperoxide Synthase 2 (PTGS2), B) Epiregulin (EREG), C) Hyaluronan Synthase 2 (HAS2) and D) TNF Alpha Induced Protein 6 (TNFAIP6) transcripts in bovine cumulus cells from COCs submitted to IVM for 3, 6, 12 or 24 hours in medium supplemented with bovine serum albumin (BSA) or fetal calf serum (FCS).




Data are the mean $\pm$ S.E.M of three independent replicates. Asterisks indicate significant differences ($P < 0.001$) between groups on a given time. Different lowercase letters indicate significant differences ($P < 0.05$) within the BSA group over time. Different uppercase letters indicate significant differences ($P < 0.05$) within the FCS group over time.

Fig. 2 – Abundance of A) X-ray repair cross complementing 6 (XRCC6), B) X-ray repair cross complementing 5 (XRCC5), C) Protein kinase DNA-activated, catalytic subunit (DNAPK), D) growth arrest and DNA damage inducible beta (GADD45B) and E) tumor protein p53 binding protein 1 (TP53BP1) transcripts in bovine cumulus cells from COCs submitted to IVM for 3, 6, 12 or 24 hours in medium supplemented with bovine serum albumin (BSA) or fetal calf serum (FCS)



Data are the mean $\pm$ S.E.M. of three independent replicates. Asterisks indicate significant differences ($P < 0.05$) between groups on a given time. Different lowercase letters indicate significant differences ($P < 0.05$) within a group over time.

Fig. 3 – Abundance of A) RAD50 double strand break repair protein (RAD50), B) RAD51 recombinase (RAD51), C) RAD52 homolog DNA repair protein (RAD52), D) ATM serine/threonine kinase (ATM), E) BRCA1 DNA repair associated (BRCA1) and F) BRCA2 DNA repair associated (BRCA2) transcripts in bovine cumulus cells from COCs submitted to IVM for 3, 6, 12 or 24 hours in medium supplemented with bovine serum albumin (BSA) or fetal calf serum (FCS).



Data are the mean $\pm$ S.E.M. of three independent replicates. Asterisks indicate significant differences ($P < 0.05$) between groups on a given time. Different lowercase letters indicate significant differences ($P < 0.05$) within a group over time.

REFERÊNCIAS

- ADAMS, R.; GIANTURCO, M. Senecio alkaloids: the alkaloids of *Senecio brasiliensis*, *fremonti* and *ambrosioides*. **Journal of the American Chemical Society**, v. 78, n. 20, p. 5315–5317, 1 out. 1956. DOI:10.1021/ja01601a044.
- AGCA, C.; YAKAN, A.; AGCA, Y. Estrus synchronization and ovarian hyper-stimulation treatments have negligible effects on cumulus oocyte complex gene expression whereas induction of ovulation causes major expression changes. **Molecular Reproduction and Development**, v. 80, n. 2, p. 102–117, 2013. DOI:10.1002/mrd.22141.
- ALI, A.; SIRARD, M.-A. Effect of the Absence or Presence of Various Protein Supplements on Further Development of Bovine Oocytes During In Vitro Maturation1. **Biology of Reproduction**, v. 66, n. 4, p. 901–905, 2002. DOI:10.1095/biolreprod66.4.901.
- ANTONIASSI, N. A. B. et al. Causas de aborto bovino diagnosticadas no Setor de Patologia Veterinária da UFRGS de 2003 a 2011. **Pesquisa Veterinária Brasileira**, v. 33, n. 2, p. 155–160, fev. 2013. DOI:10.1590/S0100-736X2013000200004.
- AUCLAIR, S. et al. Absence of cumulus cells during in vitro maturation affects lipid metabolism in bovine oocytes. **American Journal of Physiology-Endocrinology and Metabolism**, v. 304, n. 6, p. E599–E613, 2013. DOI:10.1152/ajpendo.00469.2012.
- BABAZADEH, B. et al. Protective effect of *Nigella sativa* and thymoquinone on serum/glucose deprivation-induced DNA damage in PC12 cells. **Avicenna journal of phytomedicine**, v. 2, n. 3, p. 125–32, 2012.
- BAI, P.; CANTÓ, C. The Role of PARP-1 and PARP-2 Enzymes in Metabolic Regulation and Disease. **Cell Metabolism**, v. 16, n. 3, p. 290–295, 2012. DOI:10.1016/j.cmet.2012.06.016.
- BARRETA, M. H. et al. Homologous recombination and non-homologous end-joining repair pathways in bovine embryos with different developmental competence. **Experimental Cell Research**, v. 318, n. 16, p. 2049–2058, 2012. DOI:10.1016/j.yexcr.2012.06.003.
- BARROS, C. S. L. et al. Liver biopsy for the diagnosis of *Senecio brasiliensis* (Asteraceae) poisoning in cattle. **Pesquisa Veterinária Brasileira**, v. 27, n. 1, p. 53–60, 2007. DOI:10.1590/S0100-736X2007000100010.
- BASILE, J. R. et al. Intoxicação por *Senecio* spp . (Compositae) em bovinos no sul do Brasil. **Acta Scientiae Veterinariae**, v. 33, n. 1, p. 63–68, 2005.
- BEN-AMI, I. et al. Novel function of ovarian growth factors: Combined studies by DNA microarray, biochemical and physiological approaches. **Molecular Human Reproduction**, v. 12, n. 7, p. 413–419, 2006. DOI:10.1093/molehr/gal045.

BERTOLINI, M. et al. Growth, development, and gene expression by in vivo- and in vitro-produced day 7 and 16 bovine embryos. **Molecular Reproduction and Development**, v. 63, n. 3, p. 318–328, 2002. DOI:10.1002/mrd.90015.

BIFFI, C. P. et al. Seneciosis in cattle associated with ingestion of *Senecio brasiliensis* under different forms of consumption in Santa Catarina state, Brazil. **Pesquisa Veterinaria Brasileira**, v. 39, n. 8, p. 561–563, 2019. DOI:10.1590/1678-5150-PVB-5993.

BILOTTO, S. et al. Meiosis progression and donor age affect expression profile of DNA repair genes in bovine oocytes. **Zygote**, v. 23, n. 1, p. 11–18, 2015. DOI:10.1017/S0967199413000154.

BOHRER, R. C. et al. The incidence of DNA double-strand breaks is higher in late-cleaving and less developmentally competent porcine embryos. **Biology of Reproduction**, v. 93, n. 3, p. 1–8, 2015. DOI:10.1095/biolreprod.115.130542.

BOHRER, R. C. et al. Double-strand DNA breaks are mainly repaired by the homologous recombination pathway in early developing swine embryos. **The FASEB Journal**, v. 32, n. 4, p. 1818–1829, abr. 2018. DOI:10.1096/fj.201700800R.

BORUSZEWSKA, D. et al. The effect of lysophosphatidic acid during in vitro maturation of bovine cumulus–oocyte complexes: cumulus expansion, glucose metabolism and expression of genes involved in the ovulatory cascade, oocyte and blastocyst competence. **Reproductive Biology and Endocrinology**, v. 13, n. 1, p. 44, 16 dez. 2015. DOI:10.1186/s12958-015-0044-x.

BRANZEI, D.; FOIANI, M. Regulation of DNA repair throughout the cell cycle. **Nature Reviews Molecular Cell Biology**, v. 9, n. 4, p. 297–308, 2008. DOI:10.1038/nrm2351.

BÜRKLE, A.; BENEKE, S.; MUIRAS, M. L. Poly(ADP-ribosyl)ation and aging. **Experimental Gerontology**, v. 39, n. 11–12, p. 1599–1601, 2004. DOI:10.1016/j.exger.2004.07.010.

CAIXETA, E. S. et al. Bone morphogenetic protein 15 and fibroblast growth factor 10 enhance cumulus expansion, glucose uptake, and expression of genes in the ovulatory cascade during in vitro maturation of bovine cumulus-oocyte complexes. **Reproduction**, v. 146, n. 1, p. 27–35, 2013a. DOI:10.1530/REP-13-0079.

CAIXETA, E. S. et al. Effects of FSH on the expression of receptors for oocyte-secreted factors and members of the EGF-like family during in vitro maturation in cattle. **Reproduction, Fertility and Development**, v. 25, n. 6, p. 890–899, 2013b. DOI:10.1071/RD12125.

CANDRIAN, U. et al. Transfer of orally administered 3H-seneciphylline into cow's milk. **Journal of Agricultural and Food Chemistry**, v. 39, n. 5, p. 930–933, maio 1991. DOI:10.1021/jf00005a026.

CECCALDI, R.; RONDINELLI, B.; D'ANDREA, A. D. Repair Pathway Choices and Consequences at the Double-Strand Break. **Trends in Cell Biology**, v. 26, n. 1, p. 52–64, 2016. DOI:10.1016/j.tcb.2015.07.009.

CHANG, H. et al. Effect of oocyte vitrification on DNA damage in metaphase II oocytes and the resulting preimplantation embryos. **Molecular Reproduction and Development**, v. 86, n. 11, p. 1603–1614, 2019. DOI:10.1002/mrd.23247.

CHEEKE, P. R. Toxicity and metabolism of pyrrolizidine alkaloids. **Journal of Animal Science**, v. 66, n. 9, p. 2343–2350, 1988. DOI:10.2527/jas1988.6692343x.

CHEN, L.; RUSSELL, P. T.; LARSEN, W. J. Functional Significance of Cumulus Expansion in the Mouse : Roles for the Preovulatory Synthesis of Hyaluronic Acid Within the Cumulus Mass. **Molecular Reproduction and Development**, v. 34, n. 1, p. 87–93, 1993. DOI:10.1002/mrd.1080340114.

CHENG, D. et al. The relationship between structurally different pyrrolizidine alkaloids and western flower thrips resistance in F2 hybrids of *Jacobaea vulgaris* and *Jacobaea aquatica*. **Journal of Chemical Ecology**, v. 37, n. 10, p. 1071–1080, 2011. DOI:10.1007/s10886-011-0021-6.

CICCIA, A.; ELLEDGE, S. J. The DNA Damage Response: Making It Safe to Play with Knives. **Molecular Cell**, v. 40, n. 2, p. 179–204, out. 2010. DOI:10.1016/j.molcel.2010.09.019.

CONTI, M. et al. Role of the Epidermal Growth Factor Network in Ovarian Follicles. **Molecular Endocrinology**, v. 20, n. 4, p. 715–723, 2006. DOI:10.1210/me.2005-0185.

CONTI, M. et al. Novel signaling mechanisms in the ovary during oocyte maturation and ovulation. **Molecular and Cellular Endocrinology**, v. 356, n. 1–2, p. 65–73, jun. 2012. DOI:10.1016/j.mce.2011.11.002.

DANA, E. et al. *Senecio brasiliensis* (Spreng.) Less. (Asteraceae), another potentially invasive alien species in Europe. **BioInvasions Records**, v. 10, n. 3, p. 521–536, 2021. DOI:10.3391/bir.2021.10.3.02.

DANG, Y. et al. MicroRNA-379-5p is associated with biochemical premature ovarian insufficiency through PARP1 and XRCC6. **Cell Death & Disease**, v. 9, n. 2, p. 106, 24 fev. 2018. DOI:10.1038/s41419-017-0163-8.

DAVIS, B. J. et al. Anovulation in Cyclooxygenase-2-Deficient Mice Is Restored by Prostaglandin E2 and Interleukin-1 β . **Endocrinology**, v. 140, n. 6, p. 2685–2695, 1 jun. 1999. DOI:10.1210/endo.140.6.6715.

DE VOS, M. et al. Perspectives on the development and future of oocyte IVM in clinical practice. **Journal of Assisted Reproduction and Genetics**, v. 38, n. 6, p. 1265–1280, 2021. DOI:10.1007/s10815-021-02263-5.

DEL COLLADO, M. et al. Influence of bovine serum albumin and fetal bovine serum supplementation during in vitro maturation on lipid and mitochondrial behaviour in oocytes and lipid accumulation in bovine embryos. **Reproduction, Fertility and Development**, v. 28, n. 11, p. 1721–1732, 2015. DOI:10.1071/RD15067.

DEL COLLADO, M. et al. Contributions from the ovarian follicular environment to oocyte function. **Animal Reproduction**, v. 15, n. 3, p. 261–270, 2018. DOI:10.21451/1984-3143-AR2018-0082.

DOS SANTOS, E. C. et al. YAP signaling in preovulatory granulosa cells is critical for the functioning of the EGF network during ovulation. **Molecular and Cellular Endocrinology**, v. 541, p. 111524, 2022. DOI:10.1016/j.mce.2021.111524.

DOWNS, J. A.; JACKSON, S. P. A means to a DNA end: the many roles of Ku. **Nature Reviews Molecular Cell Biology**, v. 5, n. 5, p. 367–378, 2004. DOI:10.1038/nrm1367.

EBMEYER, J. et al. Pyrrolizidine alkaloid-induced alterations of prostanoid synthesis in human endothelial cells. **Chemico-Biological Interactions**, v. 298, p. 104–111, jan. 2019. DOI:10.1016/j.cbi.2018.11.007.

FATEHI, A. N. et al. Presence of cumulus cells during in vitro fertilization protects the bovine oocyte against oxidative stress and improves first cleavage but does not affect further development. **Zygote**, v. 13, n. 2, p. 177–185, 2005. DOI:10.1017/S0967199405003126.

FENG, L. et al. Cell cycle-dependent inhibition of 53BP1 signaling by BRCA1. **Cell Discovery**, v. 1, n. 1, p. 15019, 2015. DOI:10.1038/celldisc.2015.19.

FU, P. P. et al. Pyrrolizidine alkaloids: genotoxicity, metabolism enzymes, metabolic activation and mechanisms. **Drug Metabolism Reviews**, v. 36, n. 1, p. 1–55, 26 jan. 2004. DOI:10.1081/DMR-120028426.

FÜLÖP, C. et al. Impaired cumulus mucification and female sterility in tumor necrosis factor-induced protein-6 deficient mice. **Development and Disease**, v. 130, n. 10, p. 2253–2261, 2003. DOI:10.1242/dev.00422.

GIARETTA, P. R. et al. Seneciosis in cattle associated with photosensitization. **Pesquisa Veterinária Brasileira**, v. 34, n. 5, p. 427–432, 2014. DOI:10.1590/S0100-736X2014000500007.

GORDON, G. J.; COLEMAN, W. B.; GRISHAM, J. W. Induction of cytochrome P450 enzymes in the livers of rats treated with the pyrrolizidine alkaloid retrorsine. **Experimental and Molecular Pathology**, v. 69, n. 1, p. 17–26, ago. 2000. DOI:10.1006/exmp.2000.2308.

GÓRNIK, S. L. Plantas tóxicas de interesse agropecuário. In: SPINOSA, H. S.; GÓRNIK, S. L.; PALERMO-NETO, J. (Ed.). **Toxicologia aplicada à medicina veterinária**. Barueri, SP: Manole, 2008. p. 415–458.

GRECCO, F. B. et al. Epidemiological aspects and hepatic lesions pattern in 35 outbreaks of *Senecio* spp. poisoning in cattle in southern Brazil. **Pesquisa Veterinária Brasileira**, v. 30, n. 5, p. 389–397, 2010. DOI:10.1590/S0100-736X2010000500003.

GUDMUNDSDOTTIR, K.; ASHWORTH, A. The roles of BRCA1 and BRCA2 and associated proteins in the maintenance of genomic stability. **Oncogene**, v. 25, n. 43, p. 5864–5874, 2006. DOI:10.1038/sj.onc.1209874.

GUTIÉRREZ, C. G.; CAMPBELL, B. K.; WEBB, R. Development of a long-term bovine granulosa cell culture system: induction and maintenance of estradiol production, response to follicle-stimulating hormone, and morphological characteristics. **Biology of Reproduction**, v. 56, n. 3, p. 608–616, 1997. DOI:10.1095/biolreprod56.3.608.

HARTMANN, T.; WITTE, L. Chemistry, Biology and Chemoecology of the Pyrrolizidine Alkaloids. In: PELLETIER, S. W. (Ed.). **Alkaloids: Chemical and Biological Perspectives**. Oxford: Pergamon, 1995. 9p. 155–233.

HESSEL-PRAS, S. et al. The pyrrolizidine alkaloid senecionine induces CYP-dependent destruction of sinusoidal endothelial cells and cholestasis in mice. **Archives of Toxicology**, v. 94, n. 1, p. 219–229, 2020. DOI:10.1007/s00204-019-02582-8.

HIND, D. J. N. A Checklist of the Brazilian Senecioneae (Compositae). **Kew Bulletin**, v. 48, n. 2, p. 279–295, 1993. DOI:10.2307/4117934.

HOFFMAN, B.; LIEBERMANN, D. A. Role of gadd45 in myeloid cells in response to hematopoietic stress. **Blood Cells, Molecules, and Diseases**, v. 39, n. 3, p. 344–347, 2007. DOI:10.1016/j.bcmd.2007.06.011.

HOLM, P. et al. High bovine blastocyst development in a static in vitro production system using sofaa medium supplemented with sodium citrate and myo-inositol with or without serum-proteins. **Theriogenology**, v. 52, n. 4, p. 683–700, set. 1999. DOI:10.1016/S0093-691X(99)00162-4.

HSIEH, M. et al. Luteinizing Hormone-dependent activation of the epidermal growth factor network is essential for ovulation. **Molecular and Cellular Biology**, v. 27, n. 5, p. 1914–1924, mar. 2007. DOI:10.1128/MCB.01919-06.

HUYEN, Y. et al. Methylated lysine 79 of histone H3 targets 53BP1 to DNA double-strand breaks. **Nature**, v. 432, n. 7015, p. 406–411, 2004. DOI:10.1038/nature03114.

IKEDA, S.; IMAI, H.; YAMADA, M. Apoptosis in cumulus cells during in vitro maturation of bovine cumulus-enclosed oocytes. **Reproduction**, v. 125, p. 369–376, 2003. DOI:10.1530/rep.0.1250369.

INSTITUTO BRASILEIRO DE GEOGRAFIA E ESTATÍSTICA. **Pesquisa Pecuária Municipal 2020**. Disponível em: <<https://sidra.ibge.gov.br/tabela/3939>>. Acesso em: 20 fev. 2022.

JACKSON, S. P.; BARTEK, J. The DNA-damage response in human biology and disease. **Nature**, v. 461, n. 7267, p. 1071–1078, 2009. DOI:10.1038/nature08467.

JAROUDI, S. et al. Expression profiling of DNA repair genes in human oocytes and blastocysts using microarrays. **Human Reproduction**, v. 24, n. 10, p. 2649–2655, 2009. DOI:10.1093/humrep/dep224.

JOO, J. et al. Trichosanthes kirilowii extract enhances repair of UVB radiation-induced DNA damage by regulating BMAL1 and miR-142-3p in human keratinocytes. **Molecular Medicine Reports**, v. 17, p. 877–833, 2018. DOI:10.3892/mmr.2017.7932.

KARAM, F. C.; SCHILD, A. L.; MELLO, J. R. B. Poisoning by Senecio spp. in cattle in southern Brazil: favorable conditions and control measures. **Pesquisa Veterinária Brasileira**, v. 31, n. 7, p. 603–609, 2011. DOI:10.1590/S0100-736X2011000700010.

KARAM, F. S. C. et al. Phenology of four poisonous Senecio (Asteraceae) species in southern Rio Grande do Sul, Brazil. **Pesquisa Veterinária Brasileira**, v. 22, n. 1, p. 33–39, 2002.

KARAM, F. S. C. et al. Epidemiological aspects of seneciosis in southern Rio Grande do Sul, Brazil. **Pesquisa Veterinária Brasileira**, v. 24, n. 4, p. 191–198, 2004.

KHOKHLOVA, E. V. et al. Features of DNA Repair in the Early Stages of Mammalian Embryonic Development. **Genes**, v. 11, n. 10, p. 1138, 27 set. 2020. DOI:10.3390/genes11101138.

KIMESWENGER, S. et al. Infrared A radiation promotes survival of human melanocytes carrying ultraviolet radiation-induced DNA damage. **Experimental Dermatology**, v. 25, n. 6, p. 447–452, jun. 2016. DOI:10.1111/exd.12968.

KOBAYASHI, K.; YAMASHITA, S.; HOSHI, H. Influence of epidermal growth factor and transforming growth factor- α on in vitro maturation of cumulus cell-enclosed bovine oocytes in a defined medium. **Reproduction**, v. 100, n. 2, p. 439–446, 1 mar. 1994. DOI:10.1530/jrf.0.1000439.

KOCH, J. et al. The Hippo pathway effectors YAP and TAZ interact with EGF-like signaling to regulate expansion-related events in bovine cumulus cells in vitro. **Journal of Assisted Reproduction and Genetics**, 2022. DOI:10.1007/s10815-021-02384-x.

KRABBE, A. A. et al. Megalocitose de células luteínicas grandes de vacas prenhes com seneciose crônica. **Pesquisa Veterinária Brasileira**, v. 35, n. 1, p. 33–38, 2015. DOI:10.1590/S0100-736X2015000100008.

LEIBFRIED, L.; FIRST, N. L. Characterization of bovine follicular oocytes and their ability to mature. **Journal of Animal Science**, v. 48, n. 1, p. 76–86, 1979. DOI:10.2527/jas1979.48176x.

LEIGHTON, J. K. et al. Expression of cytochrome P450 1A1, an estrogen hydroxylase, in ovarian granulosa cells is developmentally regulated. **The Journal of Steroid Biochemistry and Molecular Biology**, v. 52, n. 4, p. 351–356, 1995. DOI:10.1016/0960-0760(94)00185-O.

LEROY, J. L. et al. Comparison of Embryo Quality in High-Yielding Dairy Cows, in Dairy Heifers and in Beef Cows. **Theriogenology**, v. 64, n. 9, p. 2022–2036, 2005. DOI:10.1016/j.theriogenology.2005.05.003.

LIEBER, M. R. et al. Nonhomologous DNA end joining (NHEJ) and chromosomal translocations in humans. **Subcell. Biochem.**, v. 50, p. 279–296, 2010.

LIEBER, M. R. The Mechanism of Double-Strand DNA Break Repair by the Nonhomologous DNA End-Joining Pathway. **Annual Review of Biochemistry**, v. 79, n. 1, p. 181–211, 7 jun. 2010. DOI:10.1146/annurev.biochem.052308.093131.

LONERGAN, P. et al. Oocyte and Embryo Quality: Effect of Origin, Culture Conditions and Gene Expression Patterns. **Reproduction in Domestic Animals**, v. 38, n. 4, p. 259–267, ago. 2003. DOI:10.1046/j.1439-0531.2003.00437.x.

LORD, C. J.; ASHWORTH, A. The DNA damage response and cancer therapy. **Nature**, v. 481, n. 7381, p. 287–294, 18 jan. 2012. DOI:10.1038/nature10760.

LUCENA, R. B. et al. Poisoning by pyrrolizidine alkaloids in ruminants and horses in Brazil. **Pesquisa Veterinária Brasileira**, v. 30, n. 5, p. 447–452, 2010. DOI:10.1590/S0100-736X2010000500013.

MACHADO, M. F. et al. Fibroblast growth factor 17 and bone morphogenetic protein 15 enhance cumulus expansion and improve quality of in vitro-produced embryos in cattle. **Theriogenology**, v. 84, n. 3, p. 390–398, 2015. DOI:10.1016/j.theriogenology.2015.03.031.

MAGNUSON, B.; BEDI, K.; LJUNGMAN, M. Genome stability versus transcript diversity. **DNA Repair**, v. 44, p. 81–86, ago. 2016. DOI:10.1016/j.dnarep.2016.05.010.

MARANGOS, P. et al. DNA damage-induced metaphase I arrest is mediated by the spindle assembly checkpoint and maternal age. **Nature Communications**, v. 6, n. 1, p. 8706, 2 dez. 2015. DOI:10.1038/ncomms9706.

MARÉCHAL, A.; ZOU, L. DNA Damage Sensing by the ATM and ATR Kinases. **Cold Spring Harbor Perspectives in Biology**, v. 5, n. 9, p. a012716–a012716, 2013. DOI:10.1101/cshperspect.a012716.

MARINI, F. et al. Regulation of DNA double strand breaks processing: focus on barriers. **Frontiers in Molecular Biosciences**, v. 6, 16 jul. 2019. DOI:10.3389/fmolb.2019.00055.

MATOBA, S. et al. Predictive value of bovine follicular components as markers of oocyte developmental potential. **Reproduction, Fertility and Development**, v. 26, n. 2, p. 337, 2014. DOI:10.1071/RD13007.

MÉNDEZ, M. C.; RIET-CORREA, F. **Plantas tóxicas e micotoxícoses**. 2. ed. Pelotas, RS: Editora e Gráfica Universitária, 2008.

MOLYNEUX, R. J. et al. Pyrrolizidine alkaloid toxicity in livestock: a paradigm for human poisoning? **Food Additives & Contaminants: Part A**, v. 28, n. 3, p. 293–307, 2011. DOI:10.1080/19440049.2010.547519.

MOLYNEUX, R. J.; JOHNSON, A. E.; STUART, L. D. Delayed manifestation of senecio-induced pyrrolizidine alkaloidosis in cattle: case reports. **Veterinary and Human Toxicology**, v. 30, p. 201, 1988.

MORTON, K. M. et al. Altered mRNA expression patterns in bovine blastocysts after fertilisation in vitro using flow-cytometrically sex-sorted sperm. **Molecular Reproduction and Development**, v. 74, n. 8, p. 931–940, ago. 2007. DOI:10.1002/mrd.20573.

NAGYOVA, E. et al. Synthesis of Tumor Necrosis Factor Alpha-Induced Protein 6 in Porcine Preovulatory Follicles : A Study with A38 Antibody. **Biology of Reproduction**, v. 78, n. 5, p. 903–909, 2008. DOI:10.1095/biolreprod.107.064832.

NASCIMENTO, R. P. et al. Involvement of astrocytic CYP1A1 isoform in the metabolism and toxicity of the alkaloid pyrrolizidine monocrotaline. **Toxicon**, v. 134, p. 41–49, ago. 2017. DOI:10.1016/j.toxicon.2017.04.005.

OUANDAOGO, Z. G. et al. Differences in transcriptomic profiles of human cumulus cells isolated from oocytes at GV, MI and MII stages after in vivo and in vitro oocyte maturation. **Human Reproduction**, v. 27, n. 8, p. 2438–2447, ago. 2012. DOI:10.1093/humrep/des172.

PANIER, S.; BOULTON, S. J. Double-strand break repair: 53BP1 comes into focus. **Nature Reviews Molecular Cell Biology**, v. 15, n. 1, p. 7–18, 2014. DOI:10.1038/nrm3719.

PANIGONE, S. et al. Luteinizing Hormone Signaling in Preovulatory Follicles Involves Early Activation of the Epidermal Growth Factor Receptor Pathway. **Molecular Endocrinology**, v. 22, n. 4, p. 924–936, abr. 2008. DOI:10.1210/me.2007-0246.

PARK, J.-Y. et al. EGF-Like growth factors as mediators of LH action in the ovulatory follicle. **Science**, v. 303, n. 5658, p. 682–684, 2004. DOI:10.1126/science.1092463.

PARRISH, J. J. et al. Capacitation of bovine sperm by heparin. **Biology of Reproduction**, v. 38, n. 5, p. 1171–1180, 1 jun. 1988. DOI:10.1095/biolreprod38.5.1171.

PEDROSO, P. M. O. et al. Spontaneous poisoning in ruminants by consumption of toxic plants: cases diagnosed by the Laboratory of Veterinary Pathology (UFRGS) in the period 1996-2005. **Acta Scientiae Veterinariae**, v. 35, n. 2, p. 213–218, 2007.

PESSOA, C. R. M.; MEDEIROS, R. M. T.; RIET-CORREA, F. Economic impact, epidemiology and control poisonous plants in Brazil. **Pesquisa Veterinária Brasileira**, v. 33, n. 6, p. 752–758, 2013. DOI:10.1590/S0100-736X2013000600011.

PETERS, M. W.; PURSLEY, J. R.; SMITH, G. W. Inhibition of intrafollicular PGE2 synthesis and ovulation following ultrasound-mediated intrafollicular injection of the selective cyclooxygenase-2 inhibitor NS-398 in cattle. **Journal of Animal Science**, v. 82, n. 6, p. 1656–1662, 1 jun. 2004. DOI:10.2527/2004.8261656x.

PFAFFL, M. W. A new mathematical model for relative quantification in real-time RT-PCR. **Nucleic Acids Research**, v. 29, n. 9, p. 2002–2007, 2001. DOI:10.1093/nar/29.9.e45.

PORTELA, V. M. et al. Role of Angiotensin II in the Perioovulatory Epidermal Growth Factor-Like Cascade in Bovine Granulosa Cells In Vitro1. **Biology of Reproduction**, v. 85, n. 6, p. 1167–1174, 2011. DOI:10.1095/biolreprod.111.094193.

PRAKASH, A. S. et al. Pyrrolizidine alkaloids in human diet. **Mutation Research/Genetic Toxicology and Environmental Mutagenesis**, v. 443, n. 1, p. 53–67, 1999. DOI:10.1016/S1383-5742(99)00010-1.

RAMAKERS, C. et al. Assumption-free analysis of quantitative real-time polymerase chain reaction (PCR) data. v. 339, p. 62–66, 2003. DOI:10.1016/S0304-3940(02)01423-4.

RAMOS-IBEAS, P. et al. Senescence and Apoptosis During in vitro Embryo Development in a Bovine Model. **Frontiers in Cell and Developmental Biology**, v. 8, 18 dez. 2020. DOI:10.3389/fcell.2020.619902.

RÄTY, M. et al. In vitro maturation supplements affect developmental competence of bovine cumulus–oocyte complexes and embryo quality after vitrification. **Cryobiology**, v. 63, n. 3, p. 245–255, dez. 2011. DOI:10.1016/j.cryobiol.2011.09.134.

RENKAWITZ, J.; LADEMANN, C. A.; JENTSCH, S. Mechanisms and principles of homology search during recombination. **Nature Reviews Molecular Cell Biology**, v. 15, n. 6, p. 369–383, 2014. DOI:10.1038/nrm3805.

RICHANI, D. et al. Metabolic co-dependence of the oocyte and cumulus cells: essential role in determining oocyte developmental competence. **Human Reproduction Update**, v. 27, n. 1, p. 27–47, 2021. DOI:10.1093/humupd/dmaa043.

RICHARDS, J. S. et al. Ovulation : new dimensions and new regulators of the inflammatory-like response. **Annu. Rev. Physiol**, v. 64, p. 69–92, 2002. DOI:10.1146/annurev.physiol.64.081501.131029.

RIET-CORREA, F.; MEDEIROS, R. M. T. Intoxicações por plantas em ruminantes no Brasil e no Uruguai: importância econômica, controle e riscos para a saúde pública. **Pesquisa Veterinária Brasileira**, v. 21, n. 1, p. 38–42, 2001. DOI:10.1590/S0100-736X2001000100008.

RISSI, D. R. et al. Plant and plant-associated mycotoxins poisoning in cattle in Rio Grande do Sul, Brazil: 461 cases. **Pesquisa Veterinária Brasileira**, v. 27, n. 7, p. 261–268, 2007.

RIZOS, D. et al. Bovine Embryo Culture in the Presence or Absence of Serum: Implications for Blastocyst Development, Cryotolerance, and Messenger RNA Expression. **Biology of Reproduction**, v. 68, n. 1, p. 236–243, 2003. DOI:10.1095/biolreprod.102.007799.

RUSSELL, D. L. et al. Bidirectional communication between cumulus cells and the oocyte : Old hands and new players ? **Theriogenology**, v. 86, n. 1, p. 62–68, 2016. DOI:10.1016/j.theriogenology.2016.04.019.

SALHAB, M. et al. Kinetics of gene expression and signaling in bovine cumulus cells throughout IVM in different mediums in relation to oocyte developmental competence, cumulus apoptosis and progesterone secretion. **Theriogenology**, v. 75, n. 1, p. 90–104, 2011. DOI:10.1016/j.theriogenology.2010.07.014.

SALMON, T. B. et al. Biological consequences of oxidative stress-induced DNA damage in *Saccharomyces cerevisiae*. **Nucleic Acids Research**, v. 32, n. 12, p. 3712–3723, 2004. DOI:10.1093/nar/gkh696.

SALUSTRI, A. et al. PTX3 plays a key role in the organization of the cumulus oophorus extracellular matrix and in in vivo fertilization. **Development**, v. 131, n. 7, p. 1577–1586, 1 abr. 2004. DOI:10.1242/dev.01056.

SANDINI, T. M. et al. Prenatal exposure to integerrimine N-oxide impaired the maternal care and the physical and behavioral development of offspring rats. **International Journal of Developmental Neuroscience**, v. 36, n. 1, p. 53–63, 29 ago. 2014. DOI:10.1016/j.ijdevneu.2014.05.007.

SANDINI, T. M.; SAYURI, M.; BERTO, U. Senecio brasiliensis e alcaloides pirrolizidínicos: toxicidade em animais e na saúde humana. **Revista Biotemas Biotemas**, v. 26, n. 262, p. 83–92, 2013. DOI:10.5007/2175-7925.2013v26n2p83.

SELA-ABRAMOVICH, S. et al. Disruption of Gap Junctional Communication within the Ovarian Follicle Induces Oocyte Maturation. **Endocrinology**, v. 147, n. 5, p. 2280–2286, 1 maio 2006. DOI:10.1210/en.2005-1011.

SERRANO-BENÍTEZ, A.; CORTÉS-LEDESMA, F.; RUIZ, J. F. “An end to a means”: how DNA-end structure shapes the double-strand break repair process. **Frontiers in Molecular Biosciences**, v. 6, 10 jan. 2020. DOI:10.3389/fmolb.2019.00153.

SHIMADA, M. et al. Paracrine and Autocrine Regulation of Epidermal Growth Factor-Like Factors in Cumulus Oocyte Complexes and Granulosa Cells: Key Roles for Prostaglandin Synthase 2 and Progesterone Receptor. **Molecular Endocrinology**, v. 20, n. 6, p. 1352–1365, 2006. DOI:10.1210/me.2005-0504.

SHIMIZU, I. et al. DNA Damage Response and Metabolic Disease. **Cell Metabolism**, v. 20, n. 6, p. 967–977, 2014. DOI:10.1016/j.cmet.2014.10.008.

SILVA, C. M.; BOLZAN, A. A.; HEINZMANN, B. M. Alcalóides pirrolizidínicos em espécies do gênero *Senecio*. **Quim. Nova**, v. 29, n. 5, p. 1047–1053, 2006.

SIRARD, M. A. et al. Timing of Nuclear Progression and Protein Synthesis Necessary for Meiotic Maturation of Bovine Oocytes. **Biology of Reproduction**, v. 40, n. 6, p. 1257–1263, 1 jun. 1989. DOI:10.1095/biolreprod40.6.1257.

SKILLETER, D. N.; MATTOCKS, A. R.; NEAL, G. E. Sensitivity of different phases of the cell cycle to selected hepatotoxins in cultured liver-derived (BL9L) cells. **Xenobiotica**, v. 18, n. 6, p. 699–705, 30 jan. 1988. DOI:10.3109/00498258809041708.

SUGIMURA, S.; RICHANI, D.; GILCHRIST, R. B. Follicular guidance for oocyte developmental competence. **Animal Reproduction**, v. 15, p. 721–726, 2018. DOI:10.21451/1984-3143-AR2018-0035.

SUGIURA, K. et al. Oocyte-derived BMP15 and FGFs cooperate to promote glycolysis in cumulus cells. **Development**, v. 135, n. 4, p. 786–786, 15 fev. 2008. DOI:10.1242/dev.020024.

SUN, Q.-Y.; MIAO, Y.-L.; SCHATTEN, H. Towards a new understanding on the regulation of mammalian oocyte meiosis resumption. **Cell Cycle**, v. 8, n. 17, p. 2741–2747, 2009. DOI:10.4161/cc.8.17.9471.

SUN, X. et al. The activated DNA double-strand break repair pathway in cumulus cells from aging patients may be used as a convincing predictor of poor outcomes after in vitro fertilization-embryo transfer treatment. **Plos One**, v. 13, n. 9, p. e0204524, 2018. DOI:10.1371/journal.pone.0204524.

TANGHE, S. et al. Minireview: Functions of the cumulus oophorus during oocyte maturation, ovulation, and fertilization. **Molecular Reproduction and Development**, v. 61, n. 3, p. 414–424, mar. 2002.

TESFAYE, D. et al. Gene expression profile of cumulus cells derived from cumulus - oocyte complexes matured either in vivo or in vitro. **Reproduction, Fertility and Development**, v. 21, n. 3, p. 451, 2009. DOI:10.1071/RD08190.

THATCHER, W. W. A 100-year review: historical development of female reproductive physiology in dairy cattle. **Journal of Dairy Science**, v. 100, n. 12, p. 10272–10291, dez. 2017. DOI:10.3168/jds.2017-13399.

TITUS, S. et al. Impairment of BRCA1-Related DNA Double-Strand Break Repair Leads to Ovarian Aging in Mice and Humans. **Science Translational Medicine**, v. 5, n. 172, 2013. DOI:10.1126/scitranslmed.3004925.

TOKARNIA, C. H.; DOBEREINER, J. Intoxicação experimental por *Senecio brasiliensis* (Compositae) em bovinos. **Pesquisa Veterinária Brasileira**, v. 4, n. 2, p. 39–65, 1984.

TOKARNIA, C. H.; DÖBEREINER, J.; PEIXOTO, P. V. Poisonous plants affecting livestock in Brazil. **Toxicon**, v. 40, n. 12, p. 1635–1660, dez. 2002. DOI:10.1016/S0041-0101(02)00239-8.

TOKARNIA, C. H.; DOBEREINER, J.; PEIXOTO, P. V. *Senecio* spp. In: TOKARNIA, C. H.; DOBEREINER, J.; PEIXOTO, P. V. (Ed.). **Plantas tóxicas do Brasil para animais de produção**. 2. ed. Rio de Janeiro: Helianthus, 2012. p. 177–191.

TRIGO, J. R. et al. Chemotaxonomic value of pyrrolizidine alkaloids in southern Brazil *Senecio* (Senecioneae: Asteraceae). **Biochemical Systematics and Ecology**, v. 31, n. 9, p. 1011–1022, 2003. DOI:10.1016/S0305-1978(03)00038-3.

VAN DER VALK, J. et al. Fetal bovine serum (FBS): Past – present – future. **ALTEX**, v. 35, n. 1, p. 99–118, 2018. DOI:10.14573/altex.1705101.

VANDESOMPELE, J. et al. Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. **Genome Biol**, v. 3, n. 7, p. 1–12, 2002. DOI:https://doi.org/10.1186/gb-2002-3-7-research0034.

VELHO, F. A. de B. et al. Gene Expression and In Vitro Nuclear Maturation in Bovine Cumulus Oocyte Complexes Matured in a Medium Supplemented with Bovine Fetal Serum or Bovine Serum Albumin. **Acta Scientiae Veterinariae**, v. 41, n. 1, p. 01–07, 2013.

VENKITARAMAN, A. R. Cancer Susceptibility and the Functions of BRCA1 and BRCA2. **Cell**, v. 108, n. 2, p. 171–182, 2002. DOI:10.1016/S0092-8674(02)00615-3.

VÍTOR, A. C. et al. Studying DNA double-strand break repair: An ever-growing toolbox. **Frontiers in Molecular Biosciences**, v. 7, 21 fev. 2020. DOI:10.3389/fmolb.2020.00024.

WARD, I. M. et al. 53BP1 is required for class switch recombination. **Journal of Cell Biology**, v. 165, n. 4, p. 459–464, 2004. DOI:10.1083/jcb.200403021.

YAMASHITA, Y.; HISHINUMA, M.; SHIMADA, M. Activation of PKA, p38 MAPK and ERK1/2 by gonadotropins in cumulus cells is critical for induction of EGF-like factor and TACE/ADAM17 gene expression during in vitro maturation of porcine COCs. **Journal of Ovarian Research**, v. 2, n. 1, p. 20, 2009. DOI:10.1186/1757-2215-2-20.

YANG, M. Y.; RAJAMAHENDRAN, R. Involvement of Apoptosis in the Atresia of Nonovulatory Dominant Follicle During the Bovine Estrous Cycle. **Biology of Reproduction**, v. 63, n. 5, p. 1313–1321, 2000. DOI:10.1095/biolreprod63.5.1313.

ZHANG, T. et al. The DNA damage repair protein Ku70 regulates tumor cell and hepatic carcinogenesis by interacting with FOXO4. **Pathology - Research and Practice**, v. 212, n. 3, p. 153–161, 2016. DOI:10.1016/j.prp.2015.12.012.